

**A LECTURE NOTE
ON
TH.3 – ENGINEERING
MATHEMATICS-II
SEMESTER -2**



Prepared by – Sri Asish Sharma, (Mathematics)

Mechanical Engineering

**GOVT. POLYTECHNIC,
MALKANGIRI**

Derivative

(1)

Suppose y is a function of x i.e.

$$y = f(x).$$

Derivative of y w.r.t. x is denoted as $\frac{dy}{dx}$ or $f'(x)$.Derivative by first principle or ab-initio form:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

① Suppose $f(x) = x$ then $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$$= \lim_{h \rightarrow 0} \frac{x+h-x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

② Let us take $f(x) = x^2$ then $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$$f(x+h) = (x+h)^2 \quad \left. \begin{array}{l} \lim_{x \rightarrow a} \frac{x^n - a^n}{x-a} = na^{n-1} \\ \lim_{y \rightarrow x} \frac{y^2 - x^2}{y-x} = 2(x) \end{array} \right\} \text{Where } y = x+h$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^2 + 2hx + h^2 - x^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h(2x+h)}{h} = \lim_{h \rightarrow 0} (2x+h) = 2x + 0 = 2x$$

③ Suppose $f(x) = \sqrt{x} = x^{1/2}$, $f(x+h) = (x+h)^{1/2}$

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(x+h)^{1/2} - x^{1/2}}{h} \checkmark \\ &= \lim_{h \rightarrow 0} \frac{(x+h)^{1/2} - x^{1/2}}{(x+h) - (x)} \end{aligned}$$

Let $y = x+h$ $y \rightarrow x$ as $h \rightarrow 0$

$$= \lim_{y \rightarrow x} \frac{y^{1/2} - x^{1/2}}{y - x} \checkmark$$

$$= \frac{1}{2} (x)^{\frac{1}{2} - 1} = \frac{1}{2} (x)^{-1/2} = \frac{1}{2 x^{1/2}} = \frac{1}{2\sqrt{x}} \checkmark$$

④ Suppose $f(x) = \frac{1}{x}$ $f(x+h) = \frac{1}{x+h}$

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h} \end{aligned}$$

$$= \lim_{h \rightarrow 0} \frac{x - x - h}{x(x+h) \cdot h} =$$

$$= \lim_{h \rightarrow 0} \frac{-h}{h(x)(x+h)} = \frac{-1}{x(x+h)} = \frac{-1}{x^2}$$

$$\begin{aligned} \textcircled{8} \quad f'(x) &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} \checkmark \\ &= \lim_{h \rightarrow 0} \frac{2 \cos\left(\frac{x+h+x}{2}\right) \sin\left(\frac{x+h-x}{2}\right)}{h} \checkmark \\ &= \lim_{h \rightarrow 0} \frac{\cancel{2} \cos\left(\frac{2x+h}{2}\right) \sin(h/2)}{\cancel{2} \cdot h/2} \checkmark \end{aligned}$$

$$\begin{aligned} &= \cos\left(\frac{2x+0}{2}\right) \cdot \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin(h/2)}{(h/2)} \\ &= \cos(x) \cdot \lim_{t \rightarrow 0} \frac{\sin t}{t} \\ &= \cos(x) \cdot 1 = \cos x. \end{aligned}$$

$$\begin{aligned} \textcircled{9} \quad f(x) = \cos x \quad f'(x) &= \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-\cancel{2} \sin\left(\frac{x+h+x}{2}\right) \sin\left(\frac{x+h-x}{2}\right)}{\cancel{2} \cdot h/2} \\ &= \lim_{h \rightarrow 0} -\sin\left(\frac{2x+h}{2}\right) \boxed{\lim_{h \rightarrow 0} \frac{\sin(h/2)}{(h/2)}} \text{ form } \textcircled{8}. \\ &= -\sin\left(\frac{2x}{2}\right) (1) \\ &= -\sin(x) \end{aligned}$$

Exercise find the derivative of $\sec x, \operatorname{cosec} x$
 Ans: $\rightarrow (\sec x)' = \sec x \tan x, (\operatorname{cosec} x)' = -\operatorname{cosec}(x) \cot(x).$

$$(5) f(x) = x^n$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^n - x^n}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x+h)^n - x^n}{(x+h) - x}$$

$$\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = n a^{n-1}$$

put $x+h=y$ $y \rightarrow x$ as $h \rightarrow 0$

$$= \lim_{y \rightarrow x} \frac{y^n - x^n}{y - x}$$

$$= n x^{n-1}$$

Now you can find derivative of x^3, x^4, x^5 etc easily.

$$(6) f(x) = a^x$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{a^x (a^h) - a^x}{h}$$

$$= \lim_{h \rightarrow 0} a^x \frac{(a^h - 1)}{h}$$

$$\therefore \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \log_e a$$

$$= a^x \lim_{h \rightarrow 0} \frac{a^h - 1}{h} = a^x \log_e a = a^x \ln a$$

$$\frac{a^{m+n} - a^m}{a^m} = \frac{a^m (a^n - 1)}{a^m} = a^n - 1$$

Next take $f(x) = e^x \Rightarrow f'(x) = e^x \ln e = e^x (\because \ln e = 1)$.

$$(7) f(x) = \log_e x = \ln x$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{\ln(x+h) - \ln x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \ln \left(\frac{x+h}{x} \right)$$

$$= \lim_{h \rightarrow 0} \ln \left(1 + \frac{h}{x} \right) \frac{1}{h}$$

$$= \ln \left[\lim_{h \rightarrow 0} \left(1 + \frac{h}{x} \right)^{\frac{x}{h}} \right] \frac{1}{x} = \ln e^{1/x} = \frac{1}{x} \ln e = \frac{1}{x}$$

If $f(x) = \tan x$ then $f'(x) = \lim_{h \rightarrow 0} \frac{\tan(x+h) - \tan x}{h}$ (2)

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x} \right)$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{\sin(x+h)\cos x - \cos(x+h)\sin x}{\cos x \cdot \cos(x+h)} \right)$$

$$= \lim_{h \rightarrow 0} \frac{\sin(x+h-x)}{h \cos x \cdot \cos(x+h)}$$

$$= \lim_{h \rightarrow 0} \left(\frac{\sin h}{h} \right) \cdot \lim_{h \rightarrow 0} \frac{1}{\cos x \cdot \cos(x+h)}$$

$$= (1) \cdot \frac{1}{\cos x \cdot \cos x} = \frac{1}{\cos^2 x} = \sec^2 x.$$

Ex: \rightarrow find the derivative of $\cot x$. $A \rightarrow -\cot^2 x$.

Next

- (i) $\frac{d}{dx} (\text{any constant}) = 0.$
- (ii) $\frac{d}{dx} (f(x) \pm g(x)) = \frac{d}{dx} (f(x)) \pm \frac{d}{dx} (g(x))$
- (iii) $\frac{d}{dx} (f(x)g(x)) = f(x) \frac{d}{dx} (g(x)) + \frac{d}{dx} (f(x)) \cdot g(x).$
- (iv) $\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{g(x) \frac{d}{dx} (f(x)) - f(x) \cdot \frac{d}{dx} (g(x))}{\{g(x)\}^2}$

(v) result is true for only those $x: g(x) \neq 0.$
 (vi) $\frac{d}{dx} (af(x) + bg(x)) = a \frac{d}{dx} (f(x)) + b \frac{d}{dx} (g(x)).$
 us talk about some simple exercises.

Let

(i) $\frac{d}{dx} (x^7 + \log^2 e + e^x + a^x)$. (ii) $\frac{d}{dx} (\sin^2 x)$

$$\textcircled{3} \frac{d}{dx} \left(\frac{1-\tan x}{1+\tan x} \right) = ? \quad \textcircled{4} \frac{d}{dx} (\sin x + \cos x)^2 = ?$$

$$\textcircled{5} \frac{d}{dx} (\sin x - \cos x)^2 \quad \textcircled{6} (2x+1)^2 \quad \textcircled{7} x^7 - 10x^{11} + 5 \tan x$$

Next step is to discuss chain rule

i.e. $y = \phi(t) \quad t = \psi(x)$.

then $\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$.

a) Let us take a simple example $y = (x^2+2)^3$.

put $t = x^2+2$ then $y = t^3$.

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{d}{dt} (t^3) \cdot \frac{d}{dx} (x^2+2) \\ &= 3t^2 (2x) = 3(2x) (x^2+2)^2 \end{aligned}$$

b) Suppose $y = \sin^2 x$.

put $t = \sin x \Rightarrow y = t^2$

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{d}{dt} (t^2) \cdot \frac{d}{dx} (\sin x) \\ &= 2t \cos x \\ &= 2 \sin x \cos x \end{aligned}$$

ex Differentiate $\cos^2 x, \tan^2 x, \cot^2 x, \sec^2 x, \operatorname{cosec}^2 x$.

c) $y = \log(\sec x)$ put $\sec x = t \Rightarrow y = \ln(t)$.

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{d}{dt} \log(t) \cdot \frac{d}{dx} (\sec x) \\ &= \frac{1}{t} \cdot (\sec x \tan x) \\ &= \frac{\sec x \tan x}{\sec x} = \tan x \end{aligned}$$

$$(d) \quad y = \sqrt{\sin x}$$

put $\sin x = t$ rest is easy.

$$\text{Ans: } \rightarrow \frac{\cos x}{2\sqrt{\sin x}}$$

$$(e) \quad y = e^{\sin x} \Rightarrow y = e^t$$

put $\sin x = t \Rightarrow \frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$

$$= \frac{d}{dt}(e^t) \cdot \frac{d}{dx}(\sin x)$$
$$= e^t \cos x$$
$$= e^{\sin x} \cos x$$

$$(f) \quad y = \sqrt{\sin(\sqrt{x})}$$

put $\sqrt{x} = t \Rightarrow y = \sqrt{\sin t}$

put $\sin t = v \Rightarrow y = \sqrt{v}$

$$\frac{dy}{dx} = \frac{dy}{dv} \cdot \frac{dv}{dt} \cdot \frac{dt}{dx}$$
$$= \frac{d}{dv}(\sqrt{v}) \cdot \frac{d}{dt}(\sin t) \cdot \frac{d}{dx}(\sqrt{x})$$
$$= \frac{1}{2\sqrt{v}} \cdot \cos t \cdot \frac{1}{2\sqrt{x}}$$
$$= \frac{\cos(\sqrt{x})}{4\sqrt{x} \sqrt{\sin x}}$$

Differentiate the following

①

Answer

① $2\sqrt{x} - 3e^x + 7\sin x - 8\cos x + 11$

② $\sqrt{x} \log x$ ③ $x^2 \operatorname{cosec} x$ ④ $(x^3 - 4) \tan x$

⑤ $\frac{x-7}{x-3}$ ⑥ $\frac{e^x}{\sin x}$ ⑦ $\frac{\tan x}{x^4}$ ⑧ $\frac{1 - \cos x}{1 + \sin x}$

⑨ $\frac{4}{x} + 7\cos x - 9\log x + 8\sin x - 3$

⑩ $\sqrt{x} \log x \operatorname{cosec} x$ ⑪ $x^3 \cot x \log x$

⑫ $(2x+5)^3$ ⑬ $\sqrt{ax+b}$ ⑭ $\sin 2x$

⑮ $\cos^3 x$ ⑯ $\sin^2(5x)$ ⑰ $\sin(\cos x)$

⑱ $\cos(\log x)$ ⑲ $\log(2x+3)$ ⑳ e^{x^2}

Derivative of Inverse trigonometric functions

(1)

$$\textcircled{1} \quad y = \sin^{-1} x \quad x \in [-1, 1] \quad y \in [-\pi/2, \pi/2].$$

$$\Rightarrow x = \sin y$$

$$\Rightarrow \frac{dx}{dy} = \cos y$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\frac{dx}{dy}} = \frac{1}{\cos y} = \frac{1}{\sqrt{1-\sin^2 y}} = \frac{1}{\sqrt{1-x^2}}$$

$$\textcircled{2} \quad y = \cos^{-1} x \quad x \in [-1, 1] \quad y \in [0, \pi].$$

$$\Rightarrow x = \cos y$$

$$\Rightarrow \frac{dx}{dy} = -\sin y$$

$$\Rightarrow \frac{dy}{dx} = \frac{-1}{-\sin y} = \frac{-1}{\sqrt{1-\cos^2 y}} = \frac{-1}{\sqrt{1-x^2}}$$

$$\textcircled{3} \quad y = \tan^{-1} x \quad x \in \mathbb{R} \quad y \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

$$\Rightarrow x = \tan y$$

$$\Rightarrow \frac{dx}{dy} = \sec^2 y = 1 + \tan^2 y = 1 + x^2$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{1}{1+x^2}}$$

$$\textcircled{4} \quad y = \cot^{-1} x \quad x \in \mathbb{R} \quad y \in (0, \pi).$$

$$\Rightarrow x = \cot y$$

$$\Rightarrow \frac{dx}{dy} = -\operatorname{cosec}^2 y = -(1 + \cot^2 y) = -(1 + x^2).$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{-1}{1+x^2}}$$

$$\textcircled{5} \quad y = \sec^{-1} x. \quad x \in (-\infty, -1] \cup [1, \infty). \quad y \in \left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$$

$$y \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right].$$

$$\Rightarrow x = \sec y$$

$$\Rightarrow \frac{dx}{dy} = \sec y \tan y$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sec y \tan y} = \frac{1}{\sec y \sqrt{\sec^2 y - 1}} = \frac{1}{x \sqrt{x^2 - 1}}$$

(\because $\sec y \tan y$ is positive in the required intervals)

$$\boxed{\frac{dy}{dx} = \frac{1}{|x| \sqrt{x^2 - 1}}}$$

$$\textcircled{6} \quad y = \operatorname{cosec}^{-1} x \quad x \in (-\infty, -1] \cup [1, \infty) \quad y \in \left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$$

$$\Rightarrow x = \operatorname{cosec} y$$

$$\Rightarrow \frac{dx}{dy} = -\operatorname{cosec} y \cot y$$

$$\Rightarrow \frac{dy}{dx} = \frac{-1}{(\operatorname{cosec} y) \cot(y)} = \frac{-1}{\operatorname{cosec} y \sqrt{\operatorname{cosec}^2 y - 1}} = \frac{-1}{x \sqrt{x^2 - 1}}$$

\because $\operatorname{cosec} y \cot y$ is positive in the required intervals

$$\boxed{\frac{dy}{dx} = \frac{-1}{|x| \sqrt{x^2 - 1}}}$$

ex: $\rightarrow \frac{d}{dx} \left[\frac{1 - \tan x}{1 + \tan x} \right]^{\frac{1}{2}} = \frac{-1}{\sqrt{\cos 2x} (\cos x + \sin x)}$

Sol: $\rightarrow y = \left[\frac{1 - \tan x}{1 + \tan x} \right]^{\frac{1}{2}}$

put $t = \frac{1 - \tan x}{1 + \tan x} \Rightarrow y = \sqrt{t}$

$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{d}{dt}(\sqrt{t}) \cdot \frac{d}{dx} \left(\frac{1 - \tan x}{1 + \tan x} \right)$

$= \frac{1}{2\sqrt{t}} \left[\frac{(1 + \tan x) \frac{d}{dx}(1 - \tan x) - (1 - \tan x) \frac{d}{dx}(1 + \tan x)}{(1 + \tan x)^2} \right]$

$= \frac{1}{2\sqrt{t}} \left[\frac{(1 + \tan x)(-\sec^2 x) - (1 - \tan x)(\sec^2 x)}{(1 + \tan x)^2} \right]$

$= \frac{1}{2\sqrt{t}} \left[\frac{-\sec^2 x (1 + \tan x + 1 - \tan x)}{(1 + \tan x)^2} \right]$

$= \frac{-\sec^2 x}{2\sqrt{t} (1 + \tan x)^2}$

$= \frac{-(1 + \tan^2 x)}{\sqrt{\frac{1 - \tan x}{1 + \tan x}} (1 + \tan x)^{\frac{1}{2}} (1 + \tan x)^{\frac{3}{2}}}$

$= \frac{-(1 + \tan^2 x)}{\sqrt{1 - \tan x} \sqrt{1 + \tan x} (1 + \tan x)}$

$$\begin{aligned}
&= \frac{-(1+\tan^2 x)}{\sqrt{1-\tan^2 x} (1+\tan x)} \\
&= \frac{-(1+\tan^2 x) \sqrt{1-\tan^2 x}}{(1-\tan^2 x) (1+\tan x)} \\
&= \frac{-\sqrt{1-\tan^2 x}}{\left(\frac{1-\tan^2 x}{1+\tan^2 x}\right) (1+\tan x)} \\
&= \frac{-\sqrt{1-\tan^2 x}}{(\cos 2x) \left(1 + \frac{\sin x}{\cos x}\right)} \\
&= \frac{-\sqrt{1-\frac{\sin^2 x}{\cos^2 x}}}{\cos 2x \left(\frac{\cos 2x + \sin x}{\cos x}\right)} \\
&= \frac{-\sqrt{\frac{\cos^2 x - \sin^2 x}{\cos^2 x}}}{\cos 2x \left(\frac{\cos 2x + \sin x}{\cos x}\right)} \\
&= \frac{-\sqrt{\cos 2x}}{\cos 2x \left(\frac{\cos 2x + \sin x}{\cancel{\cos x}}\right)} \\
&= \frac{-1}{\sqrt{\cos 2x} (\cos 2x + \sin x)}
\end{aligned}$$

$$(1) \quad \frac{d}{dx} \left(\ln \tan\left(\frac{\pi}{4} + \frac{x}{2}\right) \right) = ?$$

$$y = \ln \left(\tan\left(\frac{\pi}{4} + \frac{x}{2}\right) \right)$$

$$\text{put } \frac{\pi}{4} + \frac{x}{2} = v, \quad \tan\left(\frac{\pi}{4} + \frac{x}{2}\right) = t$$

$$\therefore y = \ln t \quad \text{where } t = \tan(v).$$

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dv} \cdot \frac{dv}{dx}$$

$$= \frac{d}{dt} (\ln t) \cdot \frac{d}{dv} \tan\left(\frac{\pi}{4} + \frac{x}{2}\right) \cdot \frac{d}{dx} \left(\frac{\pi}{4} + \frac{x}{2}\right)$$

$$= \frac{1}{t} \cdot \sec^2\left(\frac{\pi}{4} + \frac{x}{2}\right) \cdot \frac{1}{2}$$

$$= \frac{1 + \tan^2 v}{2t} = \frac{1 + \tan^2\left(\frac{\pi}{4} + \frac{x}{2}\right)}{2 \tan\left(\frac{\pi}{4} + \frac{x}{2}\right)}$$

$$\text{if } \frac{\pi}{4} + \frac{x}{2} = \theta \Rightarrow \frac{dy}{dx} = \frac{1 + \tan^2 \theta}{2 \tan \theta} = \operatorname{Cosec} 2\theta$$

$$\therefore \frac{dy}{dx} = \operatorname{Cosec} 2\left(\frac{\pi}{4} + \frac{x}{2}\right) = \operatorname{Cosec}(\pi/2 + x) = \sec x$$

$$\begin{aligned} \frac{d}{dx} (\sin 5x \cos 7x) &= \sin 5x \frac{d}{dx} (\cos 7x) + \frac{d}{dx} (\sin 5x) \cos 7x \\ &= \sin 5x (-7 \sin 7x) + 5 \cos 5x \sin 7x \end{aligned}$$

$$\textcircled{4} \quad \frac{d}{dx} (\sin^{-1} 2x) = ?$$

Solⁿ: $\rightarrow y = \sin^{-1} 2x.$

put $2x = v \Rightarrow y = \sin^{-1} v.$

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{dv} \cdot \frac{dv}{dx} = \frac{d}{dv} (\sin^{-1} v) \cdot \frac{d}{dx} (2x) \\ &= \frac{1}{\sqrt{1-v^2}} (2) = \frac{2}{\sqrt{1-(2x)^2}} = \frac{2}{\sqrt{1-4x^2}}. \end{aligned}$$

$$\textcircled{5} \quad \frac{d}{dx} (\tan^{-1} (\sin^2 x))$$

Solⁿ: $\rightarrow y = \tan^{-1} (\sin^2 x).$

put $\sin^2 x = v \Rightarrow \sin x = \sqrt{v} \Rightarrow y = \tan^{-1} (v).$

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{dv} \cdot \frac{dv}{dx} = \frac{d}{dv} (\tan^{-1} v) \cdot \frac{d}{dx} (v) \\ &= \frac{1}{1+v^2} \cdot \frac{d}{dx} (\sin^2 x) \\ &= \frac{1}{1+(\sin^2 x)^2} [2 \sin x \cos x] \\ &= \frac{2 \sin x \cos x}{1+\sin^4 x} \end{aligned}$$

$$\textcircled{6} \quad \text{Diff. } \tan^{-1} \left(\frac{1}{1-x+x^2} \right)$$

Solⁿ: $\rightarrow \tan^{-1} \left(\frac{1}{1-x+x^2} \right) = \tan^{-1} \left(\frac{1}{1+x^2-x} \right)$

$$= \tan^{-1} \left(\frac{1}{1+x(x-1)} \right) = \tan^{-1} \left(\frac{x-(x-1)}{1+x(x-1)} \right)$$

$$= \tan^{-1}(x) - \tan^{-1}(x-1)$$

$$\therefore \frac{dy}{dx} = \frac{d}{dx} (\tan^{-1} x) - \frac{d}{dx} \tan^{-1}(x-1)$$

$$= \frac{1}{1+x^2} - \frac{1}{1+(x-1)^2} \cdot \frac{d}{dx}(x-1)$$

$$= \frac{1}{1+x^2} - \frac{1}{1+x^2-2x+1}$$

$$= \frac{1}{x^2+1} - \frac{1}{x^2-2x+2}$$

Ex: $\rightarrow x^2 \operatorname{cosec}^{-1}\left(\frac{1}{\ln x}\right), (x \sin^{-1} x)^{15}, \tan^{-1}(\cos \sqrt{x})$

Differentiation by Substitution

Sometimes with proper substitution we can transform the given function to a simpler function in the new variable so the process of differentiation w.r.t. the new variable becomes easier. Finally we apply chain rule to obtain the derivative w.r.t. the original variable.

e.g $\cos^{-1}(4x^3-3x)$ put $x = \cos \theta \Rightarrow \theta = \cos^{-1} x$

$$\therefore y = \cos^{-1}(4 \cos^3 \theta - 3 \cos \theta) = \cos^{-1} \cos 3\theta = 3\theta$$

$$= 3 \cos^{-1} x \quad \therefore \frac{dy}{dx} = \frac{-3}{\sqrt{1-x^2}}$$

$$1) \sin^{-1} 2x \sqrt{1-x^2}$$

$$\text{put } x = \sin \theta$$

$$\begin{aligned} \therefore y &= \sin^{-1} 2 \sin \theta \sqrt{1 - \sin^2 \theta} \\ &= \sin^{-1} 2 \sin \theta \cos \theta \\ &= \sin^{-1} \sin 2\theta \\ &= 2\theta = 2 \sin^{-1} x. \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{2}{\sqrt{1-x^2}}$$

$$② \quad \sin^{-1} \left(\frac{2x}{1+x^2} \right)$$

$$\text{put } x = \tan \theta \Rightarrow \theta = \tan^{-1} x$$

$$\therefore y = \sin^{-1} \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) = \sin^{-1} (\sin 2\theta) = 2\theta$$

$$\Rightarrow y = 2 \tan^{-1} x \Rightarrow \frac{dy}{dx} = \frac{2}{1+x^2}$$

$$③ \quad \tan^{-1} \left(\frac{2x}{1-x^2} \right) \quad \text{put } x = \tan \theta$$

$$A: \rightarrow \frac{2}{1+x^2}$$

$$④ \quad \cos^{-1} \left(\frac{1-t^2}{1+t^2} \right) \quad \text{put } t = \tan \theta$$

$$A: \rightarrow \frac{2}{1+x^2}$$

(4)

$$Q: \rightarrow \cos^{-1}(2t^2 - 1)$$

$$\text{put } t = \cos \theta \Rightarrow \theta = \cos^{-1} t$$

$$y = \cos^{-1}(2\cos^2\theta - 1) = \cos^{-1}(\cos 2\theta) = 2\theta = 2\cos^{-1} t$$

$$\Rightarrow \frac{dy}{dt} = \frac{-2}{\sqrt{1-t^2}}$$

$$Q: \rightarrow \left[\left(\frac{1+t^2}{1-t^2} \right)^2 - 1 \right]^{1/2}$$

$$y = \left[\frac{(1+t^2)^2 - (1-t^2)^2}{(1-t^2)^2} \right]^{1/2}$$

$$= \left[\frac{4t^2}{(1-t^2)^2} \right]^{1/2}$$

$$= \frac{2t}{1-t^2}$$

$$\frac{dy}{dt} = 2 \left[\frac{(1-t^2) - t(-2t)}{(1-t^2)^2} \right]$$

$$= 2 \left[\frac{1-t^2+2t^2}{(1-t^2)^2} \right] = \frac{2(1+t^2)}{(1-t^2)^2}$$

Ex: (1) using logarithm

$$\text{suppose } y = f(x)^{g(x)}$$

$$\rightarrow \log y = g(x) \log [f(x)].$$

$$\text{Q: } \rightarrow \textcircled{1} y = x^x$$

$$\Rightarrow \log y = x \log x$$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = x \cdot \frac{d}{dx}(\log x) + \frac{d}{dx}(x) \log x$$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = x \cdot \frac{1}{x} + \log x$$

$$\Rightarrow \frac{dy}{dx} = y [1 + \log x] = x^x (1 + \log x).$$

$$\textcircled{2} y = x^{\sin x}$$

$$\Rightarrow \log y = \sin x \log x$$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \sin x \frac{d}{dx}(\log x) + \frac{d}{dx}(\sin x) \log x$$

$$\Rightarrow \frac{dy}{dx} = y \left[\frac{\sin x}{x} + \cos x \log x \right]$$

$$= x^{\sin x} \left[\frac{\sin x}{x} + \cos x \log x \right].$$

③ $y = 2^{2^x}$

$\Rightarrow \log y = 2^x \log 2$

$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = (\log 2) (2^x \log 2)$

$\Rightarrow \frac{dy}{dx} = (\log 2) (2^x \log 2) (2^{2^x})$

④ $y = \left(1 + \frac{1}{x}\right)^x$

$\Rightarrow \log y = x \log \left(1 + \frac{1}{x}\right)$

$\Rightarrow \log y = x \log \left(\frac{1+x}{x}\right)$

$\Rightarrow \log y = x [\log(1+x) - \log x]$

$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = x \left[\frac{1}{1+x} - \frac{1}{x} \right] + \log(1+x) - \log x$

$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = x \left[\frac{x - (1+x)}{x(1+x)} + \log \left(\frac{1+x}{x}\right) \right]$

$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \frac{-1}{1+x} + \log \left(\frac{1+x}{x}\right)$

$\Rightarrow \frac{dy}{dx} = y \left[\frac{-1}{1+x} + \log \left(\frac{1+x}{x}\right) \right]$

$$(5) \quad y = x^{1/x} + (\sin x)^x$$

$$y = y_1 + y_2$$

$$\Rightarrow \frac{dy}{dx} = \frac{dy_1}{dx} + \frac{dy_2}{dx}$$

$$y_1 = x^{1/x}$$

$$\Rightarrow \log y_1 = \frac{1}{x} \log x$$

$$\Rightarrow \frac{1}{y_1} \cdot \frac{dy_1}{dx} = \log x \left(\frac{-1}{x^2} \right) + \frac{1}{x} \cdot \frac{1}{x}$$

$$\Rightarrow \frac{1}{y_1} \cdot \frac{dy_1}{dx} = \frac{1}{x^2} (1 - \log x)$$

$$\Rightarrow \frac{dy_1}{dx} = y_1 \left[\frac{1 - \log x}{x^2} \right] = x^{1/x} \left[\frac{1 - \log x}{x^2} \right]$$

$$y_2 = (\sin x)^x$$

$$\Rightarrow \log y_2 = x \log(\sin x)$$

$$\Rightarrow \frac{1}{y_2} \cdot \frac{dy_2}{dx} = x \cdot \frac{d}{dx} (\log \sin x) + \frac{d}{dx} (x) \cdot \log(\sin x)$$

$$\Rightarrow \frac{1}{y_2} \cdot \frac{dy_2}{dx} = x \cdot \frac{\cos x}{\sin x} + \log(\sin x)$$

$$\Rightarrow \frac{dy_2}{dx} = y_2 \left[x \cdot \cot x + \log(\sin x) \right]$$

$$= (\sin x)^x \left[x \cdot \cot x + \log(\sin x) \right]$$

Exercise Set

(1) $\frac{e^{3x^2}}{\ln \sin x}$ (2) $e^{\sin x} - a^{\cos x}$

(3) $a^{x^2} b^{x^3}$ (4) $5^{\sin x^2}$ (5) $\sqrt{\sec(2x+1)}$

(6) $\sin^2 x \cos^2 x$ (7) $\tan^3 x$ (8) $\sin^4 x$

(9) $\sqrt{a^{\sqrt{x}}}$ (10) $e^{\sqrt{ax}}$ (11) $\cos^{-1} \sqrt{\frac{1+x}{2}}$

(12) $\cot^{-1} \frac{\sqrt{1-x^2}}{x}$ (13) $\sin^{-1} \left(\frac{2\sqrt{t^2-1}}{t} \right)$

(14) $\tan^{-1} \sqrt{\frac{1-t}{1+t}}$ (15) $\sec^{-1} \left(\frac{\sqrt{a^2+x^2}}{a} \right)$

(16) $x^{\sin x} + \sin x^x$ (17) $x^{\cos x} + \cos x^x$

(18) $(\sec x + \tan x)^{\cot x}$ (19) $\frac{(x+1)(x+2)^2(x+3)^3}{(x-1)(x-2)^2(x-3)^3}$

(20) $\sin^n(x^n)$ (21) $(\sin x)^x \sqrt{\sin x} (1+x^2)^{\frac{1}{2}+x}$

Parametric differentiation

x and y both are functions of a common parameter t or θ or ϕ etc.

i.e: $x = \phi(t)$ $y = \psi(t)$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

Q! \rightarrow $x = at^2$, $y = 2at$ $\frac{dy}{dx} = ?$

Sol! \rightarrow $x = at^2 \Rightarrow \frac{dx}{dt} = 2at$

$y = 2at \Rightarrow \frac{dy}{dt} = 2a$

$$\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2a}{2at} = \frac{1}{t}$$

Q! \rightarrow $x = a \cos^3 \theta$ $y = b \sin^3 \theta$

Sol! \rightarrow $x = a \cos^3 \theta \Rightarrow \frac{dx}{d\theta} = 3a \cos^2 \theta (-\sin \theta)$

$y = b \sin^3 \theta \Rightarrow \frac{dy}{d\theta} = 3b \sin^2 \theta (\cos \theta)$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{\cancel{3}b \sin^2 \theta \cos \theta}{-\cancel{3}a \cos^2 \theta \sin \theta} = -\frac{b}{a} \tan(\theta)$$

Q! \rightarrow $x = t + \sqrt{t}$

$$\frac{dx}{dt} = 1 + \frac{1}{2\sqrt{t}}$$
$$= \frac{2\sqrt{t} + 1}{2\sqrt{t}}$$

$y = t - \sqrt{t}$

$$\frac{dy}{dt} = 1 - \frac{1}{2\sqrt{t}} = \frac{2\sqrt{t} - 1}{2\sqrt{t}}$$

$$\therefore \frac{dy}{dx} = \frac{2\sqrt{t} - 1}{2\sqrt{t}} \cdot \frac{2\sqrt{t}}{2\sqrt{t} + 1} = \frac{2\sqrt{t} - 1}{2\sqrt{t} + 1}$$

$$(4) \quad x = a(\theta + \sin\theta) \quad y = a(1 + \cos\theta)$$

$$\frac{dx}{d\theta} = a(1 + \cos\theta) \quad \frac{dy}{d\theta} = -a \sin\theta$$

$$\therefore \frac{dy}{dx} = \frac{-a \sin\theta}{a(1 + \cos\theta)} = \frac{-\sin\theta}{1 + \cos\theta}$$

exercises (1) $x = a \cos\theta, y = a \sin\theta$

(2) $x = a \cos^3 t, y = a \sin^3 t$ at $t = \pi/4$.

Solⁿ: $\rightarrow x = a \cos^3 t \Rightarrow \frac{dx}{dt} = 3a \cos^2 t (-\sin t)$

$y = a \sin^3 t \Rightarrow \frac{dy}{dt} = 3a \sin^2 t (\cos t)$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{3a \sin^2 t \cos t}{3a \cos^2 t (-\sin t)} = \tan(t)$$

$$\left. \frac{dy}{dx} \right|_{t=\pi/4} = \tan(\pi/4) = 1$$

~~(2) $x = 3 \cot t - 2 \cot^3 t$~~

(3) $x = 3 \cos t - 2 \cos^3 t, y = 3 \sin t - 2 \sin^3 t$

find $\frac{dy}{dx} = ?$

Diff. w.r.t. a function

$$y = f(x) \quad z = g(x)$$

$$\frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}}$$

(Q) \sqrt{x} w.r.t. x^2

$y = \sqrt{x} \Rightarrow \frac{dy}{dx} = \frac{1}{2\sqrt{x}}$

$z = x^2 \Rightarrow \frac{dz}{dx} = 2x$

$$\frac{dy}{dz} = \frac{\frac{1}{2\sqrt{x}}}{2x} = \frac{1}{4x^{3/2}}$$

② $y = \tan^{-1} x, \quad z = \cos^{-1} x$

$$\frac{dy}{dx} = \frac{1}{1+x^2}, \quad \frac{dz}{dx} = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{dy}{dz} = \frac{dy}{dx} \bigg/ \frac{dz}{dx} = \frac{\frac{1}{1+x^2}}{\frac{-1}{\sqrt{1-x^2}}} = -\frac{\sqrt{1-x^2}}{1+x^2}$$

③ $\sin x$ w.r.t. $\cot x$.

Solⁿ: $\rightarrow y = \sin x \Rightarrow \frac{dy}{dx} = \cos x$.

$z = \cot x \Rightarrow \frac{dz}{dx} = -\operatorname{cosec}^2 x$.

$$\frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{\cos x}{-\operatorname{cosec}^2 x} = -\sin^2 x \cos x$$

④ $\sin^{-1} \left(\frac{2x}{1+x^2} \right)$ w.r.t. $\cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)$.

Solⁿ: $\rightarrow y = \sin^{-1} \left(\frac{2x}{1+x^2} \right) = \sin^{-1} \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) = \sin^{-1} (\sin 2\theta)$
 $= 2\theta = 2 \tan^{-1} x$.

$$z = \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right) = \cos^{-1} \left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right) = \cos^{-1} (\cos 2\theta)$$

 $= 2\theta = 2 \tan^{-1} x$

$$\therefore y = z \Rightarrow \frac{dy}{dz} = 1$$

⑤ $\tan^{-1} x$ w.r.t. $\tan^{-1} \sqrt{1+x^2}$.

Differentiation of Implicit function

$F(x, y) = 0$ where x and y both are independent and dependent variables may determine one or ~~two~~ ^{more} functions.

Any such function is known as explicit function.

e.g ① $x^2 + y^2 - a^2 = 0$ find $\frac{dy}{dx} = ?$

$$\text{Sol}^n \Rightarrow 2x + 2y \frac{dy}{dx} - 0 = 0 \Rightarrow 2y \frac{dy}{dx} = -2x$$

$$\Rightarrow \frac{dy}{dx} = \frac{-2x}{2y} = \frac{-x}{y}$$

② $y^3 + 3x^2y - 2x = 0$

$$\Rightarrow 3y^2 \frac{dy}{dx} + 3 \left(x^2 \frac{dy}{dx} + y(2x) \right) - 2 = 0$$

$$\Rightarrow (3y^2 + 3x^2) \frac{dy}{dx} = \cancel{2x} - 6xy$$

$$\Rightarrow \frac{dy}{dx} = \frac{2(1 - 3xy)}{3(x^2 + y^2)}$$

③ $xy^2 + x^2y + 1 = 0$

$$\Rightarrow x(2y \frac{dy}{dx}) + y^2 + x^2 \frac{dy}{dx} + y(2x) = 0$$

$$\Rightarrow (2xy + x^2) \frac{dy}{dx} = -(2xy + y^2)$$

$$\Rightarrow \frac{dy}{dx} = \frac{-y(2x + y)}{x(2y + x)}$$

$$\textcircled{4} \quad x^y = y^x \quad \text{find } \frac{dy}{dx}$$

$$\Rightarrow y \ln x = x \ln y$$

$$\Rightarrow \frac{y}{x} + \ln x \frac{dy}{dx} = \frac{x}{y} \cdot \frac{dy}{dx} + \ln y$$

$$\Rightarrow \left(\ln x - \frac{x}{y} \right) \frac{dy}{dx} = \ln y - \frac{y}{x}$$

$$\Rightarrow \left(\frac{y \ln x - x}{y} \right) \frac{dy}{dx} = \frac{x \ln y - y}{x}$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{y(x \ln y - y)}{x(y \ln x - x)}}$$

$$\textcircled{5} \quad x^y y^x = 1 \quad \text{find } \frac{dy}{dx}$$

$$\Rightarrow \ln(x^y y^x) = \ln(1) = 0$$

$$\Rightarrow \ln(x^y) + \ln(y^x) = 0$$

$$\Rightarrow y \ln x + x \ln y = 0$$

$$\Rightarrow \ln x \cdot \frac{dy}{dx} + \frac{y}{x} + \frac{x}{y} \cdot \frac{dy}{dx} + \ln y = 0$$

$$\Rightarrow \left(\ln x + \frac{x}{y} \right) \frac{dy}{dx} = - \left(\ln y + \frac{y}{x} \right) = - \left(\frac{x \ln y + y}{x} \right)$$

$$\Rightarrow \left(\frac{y \ln x + x}{y} \right) \frac{dy}{dx} = - \frac{(x \ln y + y)}{x} \Rightarrow \frac{dy}{dx} = \frac{-y}{x} \left(\frac{x \ln y + y}{y \ln x + x} \right)$$

Exercise ① $e^{xy} + y \sin x = 1$

② $x = y \ln(xy)$

③ $y^2 \cot x = x^2 \cot y$

④ $\ln \sqrt{x^2 + y^2} = \tan^{-1} \frac{y}{x}$

⑤ $e^{xy} + y \sin x = 1$

e.g

$\sin(x+y) = y \cos(x+y)$ prove $\frac{dy}{dx} = -\frac{(1+y^2)}{y^2}$

Sol

$\sin(x+y) = y \cos(x+y)$

$\Rightarrow y = \tan(x+y)$

$\Rightarrow \frac{dy}{dx} = \sec^2(x+y) \left\{ 1 + \frac{dy}{dx} \right\}$

$\Rightarrow \frac{dy}{dx} = \left\{ 1 + \tan^2(x+y) \right\} \left\{ 1 + \frac{dy}{dx} \right\}$

$\Rightarrow \frac{dy}{dx} = (1+y^2) \left(1 + \frac{dy}{dx} \right)$

$\Rightarrow \left\{ 1 - (1+y^2) \right\} \frac{dy}{dx} = (1+y^2)$

$\Rightarrow -y^2 \frac{dy}{dx} = 1+y^2$

$\Rightarrow \frac{dy}{dx} = -\frac{(1+y^2)}{y^2}$

$$Q: \rightarrow \sqrt{1-x^4} + \sqrt{1-y^4} = k(x^2-y^2) \text{ prove } \frac{dy}{dx} = \frac{x\sqrt{1-y^4}}{y\sqrt{1-x^4}}$$

$$\text{Sol.}^n \rightarrow \text{put } x^2 = \cos\theta, y^2 = \cos\phi$$

$$\therefore \sqrt{1-\cos^2\theta} + \sqrt{1-\cos^2\phi} = k(\cos\theta - \cos\phi)$$

$$\Rightarrow \sin\theta + \sin\phi = k(\cos\theta - \cos\phi)$$

$$\Rightarrow \cancel{\sin\left(\frac{\theta+\phi}{2}\right)} \cos\left(\frac{\theta-\phi}{2}\right) = -k \left(\cancel{\sin\left(\frac{\theta+\phi}{2}\right)} \sin\left(\frac{\theta-\phi}{2}\right) \right)$$

$$\Rightarrow \tan\left(\frac{\theta-\phi}{2}\right) = \frac{-1}{k}$$

$$\Rightarrow \tan\left(\frac{\phi-\theta}{2}\right) = \frac{1}{k}$$

$$\Rightarrow \frac{1}{2}(\phi-\theta) = \tan^{-1}\left(\frac{1}{k}\right)$$

$$\Rightarrow \phi-\theta = 2 \tan^{-1}\left(\frac{1}{k}\right)$$

$$\Rightarrow \cos^{-1}(y^2) - \cos^{-1}(x^2) = 2 \tan^{-1}\left(\frac{1}{k}\right)$$

Diff. both sides w.r.t. x

$$\Rightarrow \frac{-1}{\sqrt{1-y^4}} (2y) \frac{dy}{dx} + \frac{2x}{\sqrt{1-x^4}} = 0$$

$$\Rightarrow \frac{x}{\sqrt{1-x^4}} = \frac{y}{\sqrt{1-y^4}} \frac{dy}{dx}$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{x\sqrt{1-y^4}}{y\sqrt{1-x^4}}}$$

Successive Differentiation

(1)

$$y = f(x)$$

$$\frac{dy}{dx} = f'(x)$$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right)$$

$$\frac{d^3y}{dx^3} = \frac{d}{dx} \left(\frac{d^2y}{dx^2} \right) \text{ and so on.}$$

We can write $y_1 = \frac{dy}{dx}$, $y_2 = \frac{d^2y}{dx^2}$ and so on.

ex ① $y = x^5 + 4x^3 - 2x^2 + 1$. $y_2 = ?$

Solⁿ: $\rightarrow y_1 = 5x^4 + 12x^2 - 4x$

$y_2 = 20x^3 + 24x - 4$

② $x = \sin t$, $y = \sin(pt)$ prove that

$$(1-x^2)y_2 - xy_1 + p^2y = 0$$

Solⁿ: $\rightarrow x = \sin t \Rightarrow t = \sin^{-1}x$

$$y = \sin(p \sin^{-1}x)$$

$$\Rightarrow y_1 = \cos(p \sin^{-1}x) \cdot \frac{p}{\sqrt{1-x^2}}$$

$$\Rightarrow y_1 \sqrt{1-x^2} = p \cos(p \sin^{-1}x)$$

$$\Rightarrow y_1^2 (1-x^2) = p^2 \cos^2(p \sin^{-1}x)$$

$$\Rightarrow y_1^2 (1-x^2) = p^2 (1-y^2)$$

$$y_1^2(-2x) + (1-x^2)2y_1y_2 = -1^2(-2x)$$

$$\Rightarrow (1-x^2)y_2 - xy_1 = -x^2y_1$$

$$\Rightarrow \boxed{(1-x^2)y_2 - xy_1 + xy_1 = 0}$$

③ $y = \tan^{-1}x$ Prove that $(1+x^2)y_2 + 2xy_1 = 0$

Solⁿ: $\rightarrow y = \tan^{-1}x$

$$\Rightarrow y_1 = \frac{1}{1+x^2}$$

$$\Rightarrow (1+x^2)y_1 = 1$$

Diff. both sides w.r.t. x

$$\Rightarrow (1+x^2)y_2 + 2xy_1 = 0$$

④ $2y = x\left(1 + \frac{dy}{dx}\right)$ Prove that y_2 is constant.

Solⁿ: $\rightarrow 2y = x(1+y_1)$

$$\Rightarrow 2y_1 = 1 + xy_2 + y_1$$

$$\Rightarrow y_1 = 1 + xy_2$$

$$\Rightarrow y_2 = xy_3 + y_2$$

$$\Rightarrow xy_3 = 0$$

$$\Rightarrow y_3 = 0$$

$$y_3 = \frac{d}{dx}(y_2) \therefore \boxed{y_2 = 0}$$

$$(5) \quad y = ax \sin x \quad \text{put that } x^2 y_2 - 2xy_1 + (x^2 + 2)y = 0 \quad (1)$$

$$\text{Sol}^n \rightarrow y = ax \sin x$$

$$\Rightarrow y_1 = a [x \cos x + \sin x] \quad (1)$$

$$\Rightarrow y_2 = a [x(-\sin x) + \cos x + \cos x] \quad (2)$$

$$\Rightarrow y_2 = -ax \sin x + 2a \cos x \quad (2)$$

$$\underline{\text{LHS}} \quad x^2 y_2 - 2xy_1 + (x^2 + 2)y$$

$$= x^2 (-ax \sin x + 2a \cos x) - 2x (ax \cos x + a \sin x) + (x^2 + 2)(ax \sin x)$$

$$= -a^2 x^3 \sin x + 2a^2 x^2 \cos x - 2a^2 x^2 \cos x - 2a^2 x \sin x + a^2 x^3 \sin x + 2a^2 x \sin x$$

$$= 0 \quad (\text{LHS}) \quad \text{proved}$$

$$(6) \quad y = e^{m \cos^{-1} x} \quad \text{put that } (1-x^2)y_2 - xy_1 = m^2 y$$

$$\text{Sol}^n \rightarrow y = e^{m \cos^{-1} x}$$

$$\Rightarrow y_1 = e^{\frac{-m}{\sqrt{1-x^2}}}$$

$$\Rightarrow y_1 \sqrt{1-x^2} = -m y$$

$$\Rightarrow y_1^2 (1-x^2) = m^2 y^2$$

Diff. both sides w.r.t. x

$$y_1^2 (-2x) + (1-x^2)(2y_1 y_2) = 2m^2 y_1 y_2$$

$$\Rightarrow -xy_1 + (1-x^2)y_2 = m^2 y$$

$$\Rightarrow \boxed{(1-x^2)y_2 - xy_1 = m^2 y}$$

Ex $y = e^{m \sin^{-1} x}$ prove that $(1-x^2)y_2 - xy_1 = m^2 y$

7 $y = (\sin^{-1} x)^2$ prove that $(1-x^2)y_2 - xy_1 - 2 = 0$

Solⁿ $\rightarrow y = (\sin^{-1} x)^2$

$$\Rightarrow y_1 = 2 \sin^{-1} x \left(\frac{1}{\sqrt{1-x^2}} \right)$$

$$\Rightarrow y_1 \sqrt{1-x^2} = 2 \sin^{-1} x$$

$$\Rightarrow y_1^2 (1-x^2) = 4 (\sin^{-1} x)^2$$

$$\Rightarrow y_1^2 (1-x^2) = 4y_1$$

$$\Rightarrow (1-x^2)(2y_1 y_2) - 2xy_1^2 = 4y_1$$

$$\Rightarrow (1-x^2)y_2 - xy_1 = 2$$

$$\Rightarrow \boxed{(1-x^2)y_2 - xy_1 - 2 = 0}$$

Partial Differentiation

(3)

Suppose $Z = f(x, y)$

$\frac{\partial Z}{\partial x}$ is called partial derivative of Z w.r.t. x

$\frac{\partial Z}{\partial y}$ " " " " " " " " " " " "

$$\frac{\partial Z}{\partial x} = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

$$\frac{\partial Z}{\partial y} = \lim_{k \rightarrow 0} \frac{f(x, y+k) - f(x, y)}{k}$$

$\frac{\partial Z}{\partial x}$ means Differentiation of Z w.r.t. x keeping other variables constant.

e.g.

$Z = x^2 + y^2$
 $\frac{\partial Z}{\partial x} = 2x$

$\frac{\partial Z}{\partial y} = 2y$

$$\frac{\partial^2 Z}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial Z}{\partial x} \right) \quad \frac{\partial^2 Z}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial Z}{\partial y} \right)$$

$$\frac{\partial^2 Z}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial Z}{\partial x} \right) \quad \frac{\partial^2 Z}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial Z}{\partial y} \right)$$

$$(2) z = \tan^{-1}\left(\frac{x}{y}\right).$$

$$\frac{\partial z}{\partial x} = \frac{1}{1 + \frac{x^2}{y^2}} \left(\frac{1}{y}\right)$$

$$= \frac{y^2}{y^2 + x^2} \cdot \frac{1}{y} = \frac{y}{x^2 + y^2}$$

$$\frac{\partial z}{\partial y} = \frac{1}{1 + \frac{x^2}{y^2}} \left(-\frac{x}{y^2}\right)$$

$$= \frac{-x}{x^2 + y^2}$$

$$(3) z = \ln(xy)$$

$$\frac{\partial z}{\partial x} = \frac{1}{xy} (y) = \frac{1}{x}$$

$$\frac{\partial z}{\partial y} = \frac{1}{xy} (x) = \frac{1}{y}$$

$$(4) z = e^{ax^2 + 2hxy + by^2}$$

$$\frac{\partial z}{\partial x} = e^{ax^2 + 2hxy + by^2} (2ax + 2hy)$$

$$\frac{\partial z}{\partial y} = e^{ax^2 + 2hxy + by^2} (2by + 2hx)$$

5) $Z = \sin(x^2 + y^2)$

$$\frac{\partial Z}{\partial x} = \cos(x^2 + y^2) (2x)$$

$$\frac{\partial Z}{\partial y} = \cos(x^2 + y^2) (2y)$$

6) $Z = \ln(x^2 + y^2)$

$$\frac{\partial Z}{\partial x} = \frac{1}{x^2 + y^2} (2x)$$

$$\frac{\partial Z}{\partial y} = \frac{2y}{x^2 + y^2}$$

7) $Z = a^{\frac{x}{y}}$

$$\frac{\partial Z}{\partial x} = a^{\frac{x}{y}} (\ln a) \cdot \frac{1}{y}$$

$$\frac{\partial Z}{\partial y} = a^{\frac{x}{y}} (\ln a) \left(\frac{-x}{y^2} \right)$$

8) $Z = x^2 + 2xy + y^2$

$$\frac{\partial Z}{\partial x} = 2x + 2y$$

$$\frac{\partial^2 Z}{\partial x^2} = 2$$

$$\frac{\partial Z}{\partial y} = 2y + 2x$$

$$\frac{\partial^2 Z}{\partial y^2} = 2$$

$$\frac{\partial^2 Z}{\partial y \partial x} = 2$$

$$\frac{\partial^2 Z}{\partial x \partial y} = 2$$

$$(9) \quad u = e^{x^2+y^2+z^2}$$

$$\frac{\partial u}{\partial x} = e^{x^2+y^2+z^2} (2x)$$

$$\frac{\partial u}{\partial y} = e^{x^2+y^2+z^2} (2y)$$

$$\frac{\partial u}{\partial z} = e^{x^2+y^2+z^2} (2z)$$

$$v = e^{xyz}$$

$$\frac{\partial v}{\partial x} = e^{xyz} (yz)$$

$$\frac{\partial v}{\partial y} = e^{xyz} (zx)$$

$$\frac{\partial v}{\partial z} = e^{xyz} (xy)$$

(10)

Integration: \rightarrow The process of finding the function when its derivative is given is called integration/~~Anti derivative~~. The function found is known as integral of the given function.

Simple Integration formulae

suppose $\frac{d}{dx}(g(x)) = f(x)$.

$$\Rightarrow \int f(x) dx = g(x) + c.$$

since $\frac{d}{dx}(g(x) + c) = \frac{d}{dx}(g(x))$.

$$\textcircled{1} \int x^n dx = \frac{x^{n+1}}{n+1} + K \quad (n \neq -1)$$

$$\textcircled{2} \int \frac{dx}{x} = \ln|x| + K$$

$$\textcircled{3} \int \cos x dx = \sin x + K.$$

$$\textcircled{4} \int \sin x dx = -\cos x + K$$

$$\textcircled{5} \int \sec^2 x dx = \tan x + K$$

$$\textcircled{6} \int \csc^2 x dx = -\cot x + K$$

$$\int \sec x \tan x \, dx = \sec x + K$$

$$\int \csc x \cot x \, dx = -\csc x + K$$

$$\int e^x \, dx = e^x + K$$

$$\int a^x \, dx = \frac{a^x}{\ln a} + K$$

$$\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x + K \text{ or } -\cos^{-1} x + K$$

$$\int \frac{dx}{1+x^2} = \tan^{-1} x + K \text{ or } -\cot^{-1} x + K$$

$$\int \frac{dx}{x\sqrt{x^2-1}} = \sec^{-1} x + K \text{ or } -\operatorname{cosec}^{-1} x + K$$

Algebra of Integrals

$$\textcircled{1} \int [f(x) \pm g(x)] \, dx = \int f(x) \, dx \pm \int g(x) \, dx$$

$$\textcircled{2} \int \lambda f(x) \, dx = \lambda \int f(x) \, dx, \quad \lambda \text{ is a constant.}$$

e.g $\textcircled{1} \int (x^6 + x^2 + x + 1) \, dx = \int x^6 \, dx + \int x^2 \, dx + \int x \, dx + \int 1 \, dx$
 $= \frac{x^7}{7} + \frac{x^3}{3} + \frac{x^2}{2} + x + K$

$$\textcircled{2} \int \left(\sqrt{y} + \frac{1}{\sqrt{y}} + \frac{1}{y^2} + \frac{1}{y^3} \right) dy = \int \sqrt{y} \, dy + \int \frac{dy}{\sqrt{y}} + \int \frac{dy}{y^2}$$

 $+ \int \frac{dy}{y^3}$
 $= \int y^{1/2} \, dy + \int y^{-1/2} \, dy + \int y^{-2} \, dy + \int y^{-3} \, dy$
 $= \frac{2}{3} y^{3/2} + 2y^{1/2} + (-1)y^{-1} + (-2)y^{-2} + C$

(2)

$$\begin{aligned} \textcircled{3} \int \left(4 \cos x - 3e^x + \frac{2}{\sqrt{1-x^2}} \right) dx \\ &= 4 \int \cos x dx - 3 \int e^x dx + 2 \int \frac{dx}{\sqrt{1-x^2}} \\ &= 4(\sin x) - 3e^x + 2 \sin^{-1} x + C. \end{aligned}$$

$$\begin{aligned} \textcircled{4} \int 6x^3 (x+5)^2 dx \\ &= \int 6x^3 (x^2 + 10x + 25) dx \\ &= \int 6x^5 dx + \int 60x^4 dx + \int 150x^3 dx \\ &= x^6 + 12x^5 + \frac{150x^4}{4} + K = x^6 + 12x^5 + \frac{75x^4}{2} + K. \end{aligned}$$

$$\begin{aligned} \textcircled{5} \int 5 \tan^2 x dx \\ &= \int 5 (\sec^2 x - 1) dx \\ &= 5 \left[\int \sec^2 x dx - \int dx \right] \\ &= 5 [\tan x - x] + K. \end{aligned}$$

$$\begin{aligned} \textcircled{6} \int \frac{x^4 dx}{x^2+1} &= \int \left(x^2 - 1 + \frac{1}{x^2+1} \right) dx \\ &= \int x^2 dx - \int dx + \int \frac{dx}{x^2+1} \\ &= \frac{x^3}{3} - x + \tan^{-1} x + K. \end{aligned}$$

$$\textcircled{7} \int \frac{dx}{1-\cos^2 x}$$

$$= \int \frac{dx}{\sin^2 x}$$

$$= \int \operatorname{cosec}^2 x \, dx$$

$$= -\cot x + K$$

$$\textcircled{8} \int \frac{\sin x}{\cos^2 x} \, dx$$

$$= \int \tan x \sec x \, dx$$

$$= \sec x + C$$

$$\textcircled{9} \int \left(\frac{e^x + 1}{e^x} \right) dx$$

$$= \int (e^x + e^{-x}) \, dx$$

$$= \int e^x \, dx + \int e^{-x} \, dx$$

$$= e^x - e^{-x} + C$$

$$\textcircled{10} \int \sec^2 x \operatorname{cosec}^2 x \, dx$$

$$= \int \frac{dx}{\sin^2 x \cos^2 x}$$

$$= \int \left(\frac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x} \right) dx$$

$$= \int \sec^2 x \, dx + \int \operatorname{cosec}^2 x \, dx$$

$$= \tan x - \cot x + K$$

$$\textcircled{11} \int \frac{\cos 3x \cos 2x + \sin 3x \sin 2x}{1 - \cos^2 x} dx$$

$$= \int \frac{\cos(3x - 2x)}{\sin^2 x} dx$$

$$= \int \frac{\cos x}{\sin x} \cdot \frac{1}{\sin x} dx$$

$$= \int \operatorname{cosec} x \cot x dx$$

$$= -\operatorname{cosec} x + C.$$

$\textcircled{12}$ Find the unique antiderivative $F(x)$ of

$$f(x) = 2x^2 + 1 \text{ where } F(0) = -2.$$

$$\begin{aligned} \text{Sol}^n \rightarrow F(x) &= \int (2x^2 + 1) dx \\ &= 2 \int x^2 dx + \int 1 dx + K \\ &= \frac{2x^3}{3} + x + K. \end{aligned}$$

$$F(0) = K = -2$$

$$\therefore f(x) = \frac{2}{3}x^3 + x - 2$$

$\boxed{\text{Ex}}$ $\textcircled{1} \int \left(\frac{x^4 + x^3 + x^2 + x + 2}{x^2 + 1} \right) dx$ $\textcircled{2} \int \left(\sqrt{1-x^2} + \frac{x^2}{\sqrt{1-x^2}} \right) dx$

$$\textcircled{3} \int \frac{x^6 dx}{x^2 + 1}$$

Integration by Substitution

Form-I

$$\int f(g(x)) g'(x) dx$$

put $g(x) = t \Rightarrow g'(x) dx = dt$

$$\therefore \int f(t) dt = \int f(g(x)) g'(x) dx$$

Form-II

$$\int \{f(x)\}^n f'(x) dx$$

put $f(x) = t \Rightarrow f'(x) dx = dt$

$$\int t^n dt = \frac{t^{n+1}}{n+1} + C = \frac{\{f(x)\}^{n+1}}{n+1} + C$$

Form-III

$$\int \frac{f'(x)}{f(x)} dx$$

put $f(x) = t \Rightarrow f'(x) dx = dt$

$$\int \frac{dt}{t} = \ln|t| + C = \ln|f(x)| + C$$

eg. $\int (ax+b)^n dx, n \neq -1$

put $ax+b = t$
 $\Rightarrow a = \frac{dt}{dx} \Rightarrow dx = \frac{dt}{a}$

$$= \frac{1}{a} \int t^n dt = \frac{1}{a} \frac{t^{n+1}}{n+1} + C$$

$$= \frac{(ax+b)^{n+1}}{a(n+1)} + C, n \neq -1$$

(2) $\int \cos(ax+b)dx = \frac{\sin(ax+b)}{a} + C$

(3) $\int \sin(ax+b)dx = -\frac{\cos(ax+b)}{a} + C$

(4) $\int \sec^2(ax+b)dx = \frac{\tan(ax+b)}{a} + C$

(5) $\int \operatorname{cosec}^2(ax+b)dx = -\frac{\cot(ax+b)}{a} + C$

(6) $\int \sec(ax+b) \tan(ax+b)dx = \frac{\sec(ax+b)}{a} + C$

(7) $\int \operatorname{cosec}(ax+b) \cot(ax+b)dx = -\frac{\operatorname{cosec}(ax+b)}{a} + C$

(8) $\int e^{ax+b} dx = \frac{e^{ax+b}}{a} + C$

(9) $\int \frac{dx}{\sqrt{1-(ax+b)^2}} = \frac{\sin^{-1}(ax+b)}{a} + C$

(10) $\int \frac{dx}{1+(ax+b)^2} = \frac{\tan^{-1}(ax+b)}{a} + C$

ex (1) $\int \frac{dx}{ax+b} = \frac{1}{a} \int \frac{adx}{ax+b} = \frac{1}{a} \ln|ax+b| + C$

(2) $\int \cot x dx = \int \frac{\cos x dx}{\sin x} = \int \frac{d(\sin x)}{\sin x} = \ln|\sin x| + C$

(3) $\int \tan x dx = \int \frac{\sec x \tan x dx}{\sec x} = \int \frac{d(\sec x)}{\sec x} = \ln|\sec x| + C$

$$(4) \int \sec x \, dx = \int \frac{\sec x (\sec x + \tan x)}{(\sec x + \tan x)} \, dx$$

put $\sec x + \tan x = t$

$$\Rightarrow \sec x \tan x + \sec^2 x = \frac{dt}{dx}$$

$$\Rightarrow \sec x (\sec x + \tan x) \, dx = dt$$

$$= \int \frac{dt}{t} = \ln |t| + C = \ln |\sec x + \tan x| + C$$

ex prove that $\int \sec x \, dx = \ln \tan \left(\frac{x}{4} + \frac{x}{2} \right) + C$

$$(5) \int \operatorname{cosec} x \, dx = \int \frac{\operatorname{cosec} x (\operatorname{cosec} x - \cot x)}{\operatorname{cosec} x - \cot x} \, dx$$

put $\operatorname{cosec} x - \cot x = t$

$$\Rightarrow -\operatorname{cosec} x \cot x + \operatorname{cosec}^2 x = \frac{dt}{dx}$$

$$\Rightarrow \operatorname{cosec} x (\operatorname{cosec} x - \cot x) \, dx = dt$$

$$= \int \frac{dt}{t} = \ln |t| + C$$

$$= \ln |\operatorname{cosec} x - \cot x| + C$$

ex prove that $\int \operatorname{cosec} x \, dx = \ln \left| \tan \frac{x}{2} \right| + C$

$$Q: \rightarrow \int x^2 \cos(x^3) dx$$

$$\text{put } x^3 = t$$

$$\Rightarrow 3x^2 = \frac{dt}{dx} \Rightarrow x^2 dx = \frac{dt}{3}$$

$$= \frac{1}{3} \int \cos(t) dt$$

$$= \frac{1}{3} \sin(t) + C = \frac{\sin(x^3)}{3} + C.$$

$$Q: \rightarrow \int e^x \sec(e^x) \tan(e^x) dx$$

$$\text{put } e^x = t \Rightarrow e^x dx = dt$$

$$= \int \sec(t) \tan(t) dt$$

$$= \sec(t) + C$$

$$= \sec(e^x) + C.$$

$$Q: \rightarrow \int \frac{\operatorname{cosec}^2(\ln x)}{x} dx$$

$$\text{put } \ln x = t \Rightarrow \frac{dx}{x} = dt$$

$$= \int \operatorname{cosec}^2(t) dt$$

$$= -\cot(t) + C$$

$$= -\cot(\ln x) + C.$$

$$Q: \rightarrow \int e^{x^3} x^2 dx$$

$$\text{put } x^3 = t$$

$$\Rightarrow 3x^2 = \frac{dt}{dx}$$

$$\Rightarrow x^2 dx = \frac{dt}{3}$$

$$= \frac{1}{3} \int e^t dt$$

$$= \frac{1}{3} e^t + C$$

$$= \frac{e^{x^3}}{3} + C$$

$$Q: \rightarrow \textcircled{a} \int e^{2x+7} dx = \frac{e^{2x+7}}{2} + C.$$

$$\textcircled{b} \int e^{x/3} dx = 3e^{x/3} + C.$$

$$\textcircled{c} \int e^{2 \tan x} (\sec^2 x) dx$$

$$\text{put } \tan x = t$$

$$\Rightarrow \sec^2 x = \frac{dt}{dx}$$

$$\Rightarrow \sec^2 x dx = dt$$

$$= \int e^{2t} dt$$

$$= \frac{e^{2t}}{2} + C$$

$$= \frac{e^{2(\tan x)}}{2} + C$$

$$Q: \rightarrow \int \frac{e^x + e^{-x}}{e^x - e^{-x}} dx \quad (\text{gmp})$$

$$\text{Sol}^n: \rightarrow \int \frac{e^x + e^{-x}}{e^x - e^{-x}} dx$$

$$\text{put } e^x - e^{-x} = t$$

$$\Rightarrow e^x + e^{-x} = \frac{dt}{dx}$$

$$\Rightarrow (e^x + e^{-x}) dx = dt$$

$$= \int \frac{dt}{t} = \ln|t| + C = \ln|e^x - e^{-x}| + C.$$

$$Q: \rightarrow \int \frac{dx}{x [1 + (\ln x)^2]}$$

$$\text{Sol}^n: \rightarrow \int \frac{dx}{x [1 + (\ln x)^2]}$$

$$\text{put } \ln x = t$$

$$\Rightarrow \frac{1}{x} = \frac{dt}{dx}$$

$$\Rightarrow \frac{dx}{x} = dt$$

$$= \int \frac{dt}{1 + t^2}$$

$$= \tan^{-1}(t) + C$$

$$= \tan^{-1}(\ln x) + C.$$

Integration of trigonometric functions

$$\sin(mx) \cos(nx) = \frac{1}{2} [\sin(m+n)x + \sin(m-n)x].$$

$$\sin(mx) \sin(nx) = \frac{1}{2} [\cos(m-n)x - \cos(m+n)x].$$

$$\cos(mx) \cos(nx) = \frac{1}{2} [\cos(m+n)x + \cos(m-n)x].$$

$$Q: \rightarrow \textcircled{1} \int \sin 3x \cos 2x dx = \frac{1}{2} \int [\sin(5x) + \sin x] dx$$

$$= \frac{1}{2} \left[-\frac{\cos(5x)}{5} - \cos(x) \right] + C.$$

$$\textcircled{2} \int \cos 3x \cos 4x dx = \frac{1}{2} \int [\cos 7x + \cos x] dx$$

$$= \frac{1}{2} \left[\frac{\sin(7x)}{7} + \sin x \right] + K.$$

Higher powers of sine and cosine may be simplified to sum of sines and cosines using multiple angle formula.

$$\sin^2 x = \frac{1}{2} (1 - \cos 2x)$$

$$\cos^2 x = \frac{1}{2} (1 + \cos 2x)$$

$$\cos^3 x = \frac{1}{4} (3 \cos x + \cos 3x)$$

$$\sin^3 x = \frac{1}{4} (3 \sin x - \sin 3x).$$

$$\cos^4 x = \left(\frac{1 + \cos 2x}{2} \right)^2 = \frac{1}{8} (3 + 4 \cos 2x + \cos 4x).$$

$$Q: \rightarrow \int \sin^2 x dx = \int \left(\frac{1 - \cos 2x}{2} \right) dx$$

$$= \frac{1}{2} \left[\int dx - \int \cos 2x dx \right]$$

$$= \frac{1}{2} \left[x - \frac{\sin(2x)}{2} \right] + C.$$

$$Q: \rightarrow \int \cos^3 x dx = \frac{1}{4} \int (3 \cos x + \cos 3x) dx$$

$$= \frac{1}{4} \left[3 \sin x + \frac{\sin(3x)}{3} \right] + C.$$

$$Q: \rightarrow \int \sin^4 x dx = \int \left(\frac{1 - \cos 2x}{2} \right)^2 dx$$

$$= \frac{1}{4} \left[\int \{ 1 - 2 \cos 2x + \cos^2(2x) \} dx \right]$$

$$= \frac{1}{4} \left[\int dx - 2 \int \cos 2x dx + \int \frac{1 + \cos(4x)}{2} dx \right].$$

$$= \frac{1}{4} \left[x - 2 \frac{\sin(2x)}{2} + \frac{1}{2} \left\{ x + \frac{\sin(4x)}{4} \right\} \right] + K.$$

$$Q: \rightarrow \int \sin^3 x \cos^3 x dx$$

$$\text{put } \sin x = t \Rightarrow \cos x dx = dt$$

$$= \int t^3 \cos^2 x (\cos x) dx$$

$$= \int t^3 (1-t^2) dt = \int (t^3 - t^5) dt$$

$$= \frac{t^4}{4} - \frac{t^6}{6} + C$$

$$= \frac{\sin^4 x}{4} - \frac{\sin^6 x}{6} + C.$$

$$Q: \rightarrow \int \cos^5 x \, dx$$

$$= \int \cos^4 x \cdot \cos x \, dx$$

$$\text{put } \sin x = t \\ \Rightarrow \cos x \, dx = dt$$

$$= \int (1-t^2)^2 \, dt$$

$$= \int (1-2t^2+t^4) \, dt$$

$$= \int dt - 2 \int t^2 \, dt + \int t^4 \, dt$$

$$= t - 2 \frac{t^3}{3} + \frac{t^5}{5} + K$$

$$= \sin x - \frac{2}{3} \sin^3 x + \frac{\sin^5 x}{5} + K.$$

$$Q: \rightarrow \int \sin^6 x \, dx$$

$$= \int \left(\frac{1 - \cos 2x}{2} \right)^3 \, dx.$$

$$= \frac{1}{8} \int [1 - \cos^3(2x) - 3\cos 2x + 3\cos^2(2x)] \, dx$$

$$= \frac{1}{8} \left[\int dx - \int \cos^3(2x) \, dx - 3 \int \cos 2x \, dx + 3 \int \cos^2(2x) \, dx \right].$$

$$= \frac{1}{8} \left[x - \int \left(\frac{3\cos(2x) - \cos(6x)}{4} \right) \, dx - 3 \frac{\sin(2x)}{2} \right.$$

$$\left. + 3 \int \left(\frac{1 + \cos 4x}{2} \right) \, dx \right]$$

$$= \frac{1}{8} \left[x - \frac{3}{8} \sin(2x) + \frac{\sin(6x)}{24} + \frac{3}{2 \cdot 2} \cos(2x) + \frac{3x}{2} + \frac{3}{8} \sin(4x) + K \right].$$

Integration by trigonometric substitution (8)

$$\textcircled{1} \int \frac{dx}{\sqrt{a^2-x^2}} = \sin^{-1}\left(\frac{x}{a}\right) + K.$$

$$\textcircled{2} \int \frac{dx}{a^2+x^2} = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + K.$$

$$\textcircled{3} \int \frac{dx}{\sqrt{x^2+a^2}} = \ln|x+\sqrt{x^2+a^2}| + K.$$

$$\textcircled{4} \int \frac{dx}{\sqrt{x^2-a^2}} = \ln|x+\sqrt{x^2-a^2}| + K.$$

$$Q: \rightarrow \textcircled{1} \int \frac{dx}{\sqrt{25-16x^2}} = \int \frac{dx}{\sqrt{5^2-(4x)^2}} = \frac{1}{4} \sin^{-1}\left(\frac{4x}{5}\right) + C.$$

$$\textcircled{2} \int \frac{e^x dx}{e^{2x}+9} \quad \text{put } e^x = t \Rightarrow e dx = dt$$

$$= \int \frac{dt}{t^2+9} = \frac{1}{3} \tan^{-1}\left(\frac{t}{3}\right) + C = \frac{1}{3} \tan^{-1}\left(\frac{e^x}{3}\right) + C$$

$$\textcircled{3} \int \frac{\cos\theta d\theta}{\sqrt{4\sin^2\theta+1}} = \int \frac{dt}{\sqrt{1+4t^2}} \quad \text{if we put } \sin\theta = t$$

$$= \frac{1}{2} \tan^{-1}(2t) + C$$
$$= \frac{1}{2} \tan^{-1}(2\sin\theta) + C$$

$$= \frac{1}{2} \ln|t + \sqrt{1+4t^2}| + C$$

$$= \frac{1}{2} \ln|\sin\theta + \sqrt{1+4\sin^2\theta}| + C.$$

$$\textcircled{4} \int \frac{dx}{x \sqrt{x^8-4}}$$

$$= \frac{1}{4} \int \frac{4x^3 dx}{x^4 \sqrt{x^8-4}}$$

$$\text{put } x^4 = t \Rightarrow 4x^3 dx = dt$$

$$= \frac{1}{4} \int \frac{dt}{t \sqrt{t^2-2^2}}$$

$$= \frac{1}{4} \cdot \frac{1}{2} \sec^{-1}\left(\frac{t}{2}\right) + K$$

$$= \frac{1}{8} \sec^{-1}\left(\frac{x^4}{2}\right) + K.$$

$$\int \frac{(x+5) dx}{\sqrt{x^2+6x-7}}$$

$$= \int \frac{x+3+2}{\sqrt{(x+3)^2-16}} dx$$

$$= \int \frac{z+2}{\sqrt{z^2-16}} \text{ put } x+3=z$$

$$= \int \frac{z dz}{\sqrt{z^2-16}} + 2 \int \frac{dz}{\sqrt{z^2-16}}$$

Rest left as exercise.

$$Q: \rightarrow \int \frac{x+3}{\sqrt{9-x^2}} dx$$

$$= \int \frac{x dx}{\sqrt{9-x^2}} + 3 \int \frac{dx}{\sqrt{3^2-x^2}}$$

$$= \frac{-1}{2} \int \frac{-2x dx}{\sqrt{9-x^2}} + 3 \sin^{-1}\left(\frac{x}{3}\right) + K$$

$$= \frac{-1}{2} \int \frac{dv}{\sqrt{v}} \quad \begin{array}{l} \text{put } 9-x^2=v \\ \Rightarrow -2x dx = dv \end{array}$$

$$= \frac{-1}{2} \int \frac{dv}{\sqrt{v}} + 3 \sin^{-1}\left(\frac{x}{3}\right) + K$$

$$= -\sqrt{v} + 3 \sin^{-1}\left(\frac{x}{3}\right) + K$$

$$= -\sqrt{9-x^2} + 3 \sin^{-1}\left(\frac{x}{3}\right) + K$$

$$Q: \rightarrow \int \frac{(x+3) dx}{\sqrt{5-x^2-4x}}$$

$$= \int \frac{(x+3) dx}{\sqrt{5+4-4-x^2-4x}}$$

$$= \int \frac{(x+3) dx}{\sqrt{9-(x^2+4x+4)}}$$

$$= \int \frac{(x+3) dx}{\sqrt{9-(x+2)^2}}$$

$$= \int \frac{(x+2)+1}{\sqrt{9-(x+2)^2}} dx$$

$$= \int \frac{(x+2)dx}{\sqrt{3^2 - (x+2)^2}} + \int \frac{dx}{\sqrt{3^2 - (x+2)^2}}$$

put $3^2 - (x+2)^2 = v$

$$\Rightarrow -2(x+2) = \frac{dv}{dx}$$

$$\Rightarrow (x+2)dx = \frac{-dv}{2}$$

$$= \frac{-1}{2} \int \frac{dv}{\sqrt{v}} + \sin^{-1} \left(\frac{x+2}{3} \right) + K$$

$$= - \int \frac{dv}{2\sqrt{v}} + \sin^{-1} \left(\frac{x+2}{3} \right) + K$$

$$= -\sqrt{v} + \sin^{-1} \left(\frac{x+2}{3} \right) + K$$

$$= -\sqrt{9 - (x+2)^2} + \sin^{-1} \left(\frac{x+2}{3} \right) + K.$$

$$\rightarrow \int \frac{x^9}{x^{20} + 4} dx$$

$$= \int \frac{x^9 dx}{(x^{10})^2 + 2^2}$$

put $x^{10} = t$

$$\Rightarrow 10x^9 = \frac{dt}{dx}$$

$$\Rightarrow x^9 dx = \frac{dt}{10}$$

$$= \frac{1}{10} \int \frac{dt}{t^2 + 2}$$

$$= \frac{1}{10} \cdot \frac{1}{2} \tan^{-1} \left(\frac{t}{2} \right) + K$$

$$= \frac{1}{20} \tan^{-1} \left(\frac{x^{10}}{2} \right) + K$$

$$\begin{aligned}
 Q: \rightarrow & \int \frac{(x+5) dx}{x^2+6x+13} \\
 = & \int \frac{(x+5) dx}{x^2+6x+9+4} \\
 = & \int \frac{(x+5) dx}{(x+3)^2+2^2} \\
 = & \int \frac{(x+3)+2}{(x+3)^2+2^2} dx \\
 = & \int \frac{(x+3) dx}{(x+3)^2+2^2} + 2 \int \frac{dx}{(x+3)^2+2^2} \\
 = & \frac{1}{2} \int \frac{2(x+3) dx}{(x+3)^2+2^2} + \frac{2}{2} \tan^{-1} \left(\frac{x+3}{2} \right) + K \\
 = & \frac{1}{2} \int \frac{dv}{v} + \tan^{-1} \left(\frac{x+3}{2} \right) + K \\
 & \text{where } v = (x+3)^2+2^2 \Rightarrow 2(x+3) dx = dv \\
 = & \frac{1}{2} \ln |v| + \tan^{-1} \left(\frac{x+3}{2} \right) + K \\
 = & \frac{1}{2} \ln |x^2+6x+13| + \tan^{-1} \left(\frac{x+3}{2} \right) + K.
 \end{aligned}$$

$$Q: \rightarrow \int \frac{dx}{\sqrt{3x^2+4}} = \int \frac{dx}{\sqrt{(\sqrt{3}x)^2+2^2}} = \frac{1}{\sqrt{3}} \ln |\sqrt{3}x + \sqrt{3x^2+4}| + K$$

$$Q: \rightarrow \int \frac{4e^x dx}{\sqrt{3e^{2x} + 4}}$$

$$= 4 \int \frac{e^x dx}{\sqrt{(\sqrt{3}e^x)^2 + 2^2}}$$

$$\text{put } \sqrt{3}e^x = t \\ \Rightarrow \sqrt{3}e^x dx = dt$$

$$= \frac{4}{\sqrt{3}} \int \frac{dt}{\sqrt{t^2 + 4}}$$

$$= \frac{4}{\sqrt{3}} \log |t + \sqrt{t^2 + 4}| + K$$

$$= \frac{4}{\sqrt{3}} \log |\sqrt{3}e^x + \sqrt{3e^{2x} + 4}| + K$$

$$Q: \rightarrow \int \frac{x^2 dx}{\sqrt{x^6 + a^6}}$$

$$= \frac{1}{3} \int \frac{3x^2 dx}{\sqrt{(x^3)^2 + (a^3)^2}}$$

$$\text{put } x^3 = v \\ \Rightarrow 3x^2 dx = dv$$

$$= \frac{1}{3} \int \frac{dv}{\sqrt{v^2 + (a^3)^2}}$$

$$= \frac{1}{3} \log |v + \sqrt{v^2 + a^6}| + K$$

$$= \frac{1}{3} \log |x^3 + \sqrt{x^6 + a^6}| + K$$

Q: $\rightarrow \int \frac{2x+11}{\sqrt{x^2+10x+29}} dx$

$= \int \frac{2x+11}{\sqrt{x^2+10x+25+4}} dx$

$= \int \frac{2x+11}{\sqrt{(x+5)^2+2^2}} dx$

$= \int \frac{\{2(x+5)+1\} dx}{\sqrt{(x+5)^2+2^2}}$

$= \int \frac{2(x+5) dx}{\sqrt{(x+5)^2+2^2}} + \int \frac{dx}{\sqrt{(x+5)^2+2^2}}$

put $(x+5)^2+2^2 = v$
 $\Rightarrow 2(x+5) dx = dv$

$= \frac{1}{2} \int \frac{dv}{\sqrt{v}} + \ln |(x+5) + \sqrt{(x+5)^2+2^2}| + K$

$= \frac{1}{2} \cdot 2 \sqrt{v} + \ln |(x+5) + \sqrt{(x+5)^2+2^2}| + K$

$= 2 \sqrt{x^2+10x+29} + \ln |(x+5) + \sqrt{x^2+10x+29}| + K$

$$\int \frac{e^{5x} dx}{\sqrt{e^{10x} - 4}}$$

put $e^{5x} = t$
 $\Rightarrow 5e^{5x} = \frac{dt}{dx}$

$$\Rightarrow e^{5x} dx = \frac{dt}{5}$$

$$= \frac{1}{5} \int \frac{dt}{\sqrt{t^2 - 4}}$$

$$= \frac{1}{5} \ln |t + \sqrt{t^2 - 4}| + K$$

$$= \frac{1}{5} \ln |e^{5x} + \sqrt{e^{10x} - 4}| + K$$

$$\int \frac{\cos \theta d\theta}{\sin^2 \theta \sqrt{\csc^2 \theta - 4}} = \int \frac{\frac{\cos \theta}{\sin \theta} \cdot \frac{1}{\sin \theta} d\theta}{\sqrt{\csc^2 \theta - 4}}$$

~~$$= \int \frac{\cos \theta d\theta}{\sin^2 \theta \sqrt{\frac{1}{\sin^2 \theta} - 4}}$$~~

~~$$= \int \frac{\cos \theta d\theta}{\sin^2 \theta \sqrt{1 - 4\sin^2 \theta}}$$~~

~~$$= \int \frac{\cos \theta d\theta}{\sin \theta \sqrt{1 - (2\sin \theta)^2}}$$~~

~~put $2\sin \theta = t$
 $\Rightarrow 2\cos \theta = \frac{dt}{d\theta}$~~

~~$$\Rightarrow \cos \theta d\theta = \frac{dt}{2}$$~~

~~$$= \int \frac{dt}{(2t) \sqrt{1 - t^2}}$$~~

~~$$= \frac{1}{2} \int \frac{dv}{\sqrt{1 - v^2}}$$~~

$$= \int \frac{\operatorname{cosec} \theta \cot \theta d\theta}{\sqrt{\csc^2 \theta - 4}}$$

put $\operatorname{cosec} \theta = t$
 $\Rightarrow -\operatorname{cosec} \theta \cot \theta = \frac{dt}{d\theta}$

$$\Rightarrow \operatorname{cosec} \theta \cot \theta d\theta = -dt$$

$$= - \int \frac{dt}{\sqrt{t^2 - 4}}$$

$$= - \log |t + \sqrt{t^2 - 4}| + K$$

~~$$= - \log |\operatorname{cosec} \theta + \sqrt{\csc^2 \theta - 4}|$$~~

~~$$= - \log |\operatorname{cosec} \theta + \sqrt{\csc^2 \theta - 4}|$$~~

+ C

put $2t = v$
 $\Rightarrow dt = \frac{dv}{2}$

Integration by parts

If v and w are differentiable functions of

$$x \text{ then } \frac{d}{dx}(vw) = v \frac{dw}{dx} + w \frac{dv}{dx}$$

$$\Rightarrow v \frac{dw}{dx} = \frac{d}{dx}(vw) - w \frac{dv}{dx}$$

$$\Rightarrow \int v \frac{dw}{dx} dx = \int \frac{d}{dx}(vw) dx - \int w \frac{dv}{dx} dx$$

$$\text{Setting } u = \frac{dw}{dx} \Rightarrow w = \int u dx$$

$$\Rightarrow \int uv dx = v \int u dx - \int \left[\int u dx \cdot \frac{dv}{dx} \right] dx$$

$$\Rightarrow \int uv dx = v \int u dx - \int \left[\int u dx \cdot \frac{dv}{dx} \right] dx$$

$$\int (\text{Int of product}) dx = (\text{1st function}) \int (\text{2nd function}) dx - \int \left[\frac{d}{dx} (\text{1st function}) \left(\int \text{2nd function} \right) dx \right]$$

So 1st function and 2nd function is not that significant.

Take a keyword ILATE

I \rightarrow Inverse trigonometric functions

L \rightarrow Logarithm functions

A \rightarrow Algebraic function

T \rightarrow Trigonometric function

E \rightarrow Exponential function

Whichever letter comes first take that as first and other as second function.

eg) $\int (\log x) dx = \int (\log x) \cdot 1 dx$

$\log x$ is logarithmic function

1 is algebraic function

So $\log x$ is 1st and 1 is 2nd function

$$= (\log x) \int 1 dx - \int \left[\frac{d}{dx} (\log x) (\int 1 dx) \right] dx$$

$$= (\log x) (x) - \int \left(\frac{1}{x} \right) (x) dx$$

$$= x \log x - \int dx + K$$

$$= \boxed{x \log x - x + K}$$

$$Q: \rightarrow \int x \cos x \, dx$$

$$= (x) \int \cos x \, dx - \int \left[\frac{d}{dx}(x) \int \cos x \, dx \right] dx$$

$$= (x)(\sin x) - \int (1)(\sin x) \, dx$$

$$= x(\sin x) - \int \sin x \, dx$$

$$= x(\sin x) - (-\cos x) + K$$

$$= x(\sin x) + \cos x + K$$

$$Q: \rightarrow \int x e^x \, dx$$

$$= (x) \int e^x \, dx - \int \left[\frac{d}{dx}(x) \int e^x \, dx \right] dx$$

$$= (x)(e^x) - \int (1) \cdot e^x \, dx$$

$$= x e^x - \int e^x \, dx + K$$

$$= x e^x - e^x + K \quad \text{--- (1)}$$

$$Q: \rightarrow \int x^2 e^x \, dx$$

$$= (x^2) \int e^x \, dx - \int \left[\frac{d}{dx}(x^2) \int e^x \, dx \right] dx$$

$$= (x^2) \cdot e^x - \int (2x) \cdot e^x \, dx$$

$$= x^2 e^x - 2 \int x e^x \, dx$$

$$= x^2 e^x - 2 [x e^x - e^x] + K$$

$$= x^2 e^x - 2x e^x + 2e^x + K$$

Ex Verify $\int x^n e^x dx$
 $= (x^n) \cdot e^x - n(x^{n-1}) \cdot e^x + \frac{n}{2} x^{n-2} e^x + \dots + n! e^x + K.$

Q: $\rightarrow \int \tan^{-1} x dx$

$= \int (\tan^{-1} x) \cdot 1 dx$

~~(1) f~~ $= \tan^{-1}(x) \int 1 dx - \int \left[\frac{d}{dx} (\tan^{-1} x) \int 1 dx \right] dx$

$= \tan^{-1}(x) \cdot x - \int \left(\frac{1}{1+x^2} \right) (x) dx$

$= x \tan^{-1} x - \frac{1}{2} \int \frac{2x dx}{1+x^2}$

$= x \tan^{-1} x - \frac{1}{2} \log(1+x^2) + K.$

$\int e^x [f(x) + f'(x)] dx = e^x f(x) + C.$

look $\frac{d}{dx} [e^x f(x)] = e^x f'(x) + f(x) e^x$
 $= e^x (f(x) + f'(x))$

$\therefore \int e^x [f(x) + f'(x)] dx = e^x f(x) + K$

$$Q: \rightarrow \int \frac{e^x}{x} (1 + x \ln x) dx$$

$$= \int e^x \left(\frac{1}{x} + \ln x \right) dx$$

$$\text{Look } f(x) = \ln x \Rightarrow f'(x) = \frac{1}{x}$$

$$= \int e^x \left(\ln x + \frac{d}{dx} (\ln x) \right) dx$$

$$= e^x \ln x + K.$$

$$Q: \rightarrow \int e^x (\tan x + \ln \sec x) dx$$

easy take as exercise Ans: $\rightarrow e^x \ln(\sec x) + K.$

$$Q: \rightarrow \int e^x (\cot x + \ln \sin x) dx = e^x \ln(\sin x) + K.$$

$$Q: \rightarrow \int \frac{x e^x dx}{(1+x)^2}$$

$$= \int \frac{(1+x-1)e^x dx}{(1+x)^2}$$

$$= \int e^x \left\{ \frac{1}{1+x} - \frac{1}{(1+x)^2} \right\} dx$$

$$\text{put } \frac{1}{1+x} = f(x) \Rightarrow \frac{-1}{(1+x)^2} = f'(x)$$

$$\therefore \int e^x \left[\frac{1}{1+x} - \frac{1}{(1+x)^2} \right] dx$$

$$= \frac{e^x}{1+x} + K.$$

$$Q: \rightarrow \int e^x \left(\frac{1+\sin x}{1+\cos x} \right) dx$$

$$= \int e^x \left(\frac{1}{1+\cos x} + \frac{\sin x}{1+\cos x} \right) dx$$

$$\text{Let } f(x) = \frac{\sin x}{1+\cos x}$$

$$f'(x) = \frac{(1+\cos x) \cos x - \sin x (-\sin x)}{(1+\cos x)^2}$$

$$= \frac{\cos x + \cos^2 x + \sin^2 x}{(1+\cos x)^2}$$

$$= \frac{1+\cos x}{(1+\cos x)^2} = \frac{1}{1+\cos x}$$

$$\therefore \int e^x \left(\frac{1}{1+\cos x} + \frac{\sin x}{1+\cos x} \right) dx$$

$$= \frac{e^x \sin x}{1+\cos x} + K.$$

Q: $\rightarrow \int e^{ax} \cos bx \, dx = I$ (say)

$$= \cos bx \cdot \int e^{ax} \, dx - \int \left[\frac{d}{dx} (\cos bx) \cdot \int e^{ax} \, dx \right] dx$$

$$= \cos bx \cdot \frac{e^{ax}}{a} - \int (-b \sin bx) \frac{e^{ax}}{a} \, dx$$

$$= \frac{e^{ax} \cos bx}{a} + \frac{b}{a} \int e^{ax} \sin bx \, dx.$$

$$= \frac{e^{ax} \cos bx}{a} + \frac{b}{a} \left[\sin bx \int \frac{e^{ax}}{a} \, dx - \int \left[\frac{d}{dx} (\sin bx) \int e^{ax} \, dx \right] dx \right]$$

$$\frac{e^{ax} \cos bx}{a} + \frac{b}{a} \left[\frac{\sin bx \cdot e^{ax}}{a} - \int b \cos bx \cdot \frac{e^{ax}}{a} \, dx \right].$$

$$\frac{e^{ax} \cos bx}{a} + \frac{b}{a^2} e^{ax} \sin bx - \frac{b^2}{a^2} I$$

$$\Rightarrow \left(I + \frac{b^2}{a^2} I \right) = e^{ax} \left[\frac{b \sin bx + a \cos bx}{a^2} \right].$$

$$\Rightarrow \left(\frac{a^2 + b^2}{a^2} \right) I = \frac{e^{ax} (a \cos bx + b \sin bx)}{a^2}$$

$$\Rightarrow I = e^{ax} \left(\frac{a \cos bx + b \sin bx}{a^2 + b^2} \right) + K$$

Ex Find $\int e^{ax} \sin(bx) dx = ?$

Ans: $\rightarrow \frac{e^{ax}}{a^2+b^2} (a \sin bx - b \cos bx) + K.$

Formula

$$\int \sqrt{a^2-x^2} dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1}\left(\frac{x}{a}\right) + K.$$

$$\int \sqrt{a^2+x^2} dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \ln|x+\sqrt{x^2+a^2}| + K$$

$$\int \sqrt{x^2-a^2} dx = \frac{x}{2} \sqrt{x^2-a^2} - \frac{a^2}{2} \ln|x+\sqrt{x^2-a^2}| + K$$

Proof $I = \int \sqrt{a^2-x^2} dx = \sqrt{a^2-x^2} \int 1 dx - \int \left[\frac{d}{dx} (\sqrt{a^2-x^2}) \int 1 dx \right] dx$

$$= \sqrt{a^2-x^2} (x) - \int \frac{(-2x)}{2\sqrt{a^2-x^2}} \cdot x dx$$

$$= x\sqrt{a^2-x^2} - \int \frac{-x^2 dx}{\sqrt{a^2-x^2}}$$

$$= x\sqrt{a^2-x^2} - \int \frac{(a^2-x^2-a^2) dx}{\sqrt{a^2-x^2}} + a^2 \int \frac{dx}{\sqrt{a^2-x^2}}$$

$$= x\sqrt{a^2-x^2} - I + a^2 \sin^{-1}\left(\frac{x}{a}\right).$$

$$\therefore 2I = x\sqrt{a^2-x^2} + a^2 \sin^{-1}\left(\frac{x}{a}\right)$$

$$\Rightarrow \boxed{I = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1}\left(\frac{x}{a}\right)}$$

Other two parts are easy and similar to the last one. Work out them.

$$\begin{aligned}
 Q: & \rightarrow \int \ln(x + \sqrt{x^2 + a^2}) dx \\
 & = \ln(x + \sqrt{x^2 + a^2}) \int 1 dx - \int \left[\frac{d}{dx} \ln(x + \sqrt{x^2 + a^2}) \int 1 dx \right] dx \\
 & = x \ln(x + \sqrt{x^2 + a^2}) - \int \left[\frac{1}{x + \sqrt{x^2 + a^2}} \left(1 + \frac{x}{\sqrt{x^2 + a^2}} \right) \cdot x \right] dx \\
 & = x \ln(x + \sqrt{x^2 + a^2}) - \int \left(\frac{1}{x + \sqrt{x^2 + a^2}} \right) \left(\frac{x + \sqrt{x^2 + a^2}}{\sqrt{x^2 + a^2}} \right) x dx \\
 & = x \ln(x + \sqrt{x^2 + a^2}) - \frac{1}{2} \int \frac{2x dx}{\sqrt{x^2 + a^2}} \\
 & = x \ln(x + \sqrt{x^2 + a^2}) - \int \frac{dv}{2\sqrt{v}} \quad \text{put } v = x^2 + a^2 \\
 & \quad \quad \quad \Rightarrow 2x dx = dv \\
 & = x \ln(x + \sqrt{x^2 + a^2}) - \sqrt{v} + C \\
 & = x \ln(x + \sqrt{x^2 + a^2}) - \sqrt{x^2 + a^2} + C
 \end{aligned}$$

ex $\int \ln(x + \sqrt{x^2 - a^2}) dx.$

$$Q: \rightarrow ① \int \sqrt{9-x^2} dx$$

$$② \int \sqrt{5-4x^2} dx$$

Both are easy exercises. Use formula to get them.

$$③ \int \sqrt{1-x^2-2x} dx$$

$$= \int \sqrt{1-x^2-2x-1+1} dx$$

$$= \int \sqrt{2-(x^2+2x+1)} dx$$

$$= \int \sqrt{(\sqrt{2})^2 - (x+1)^2} dx$$

$$\text{put } (x+1) = t \Rightarrow dx = dt$$

$$= \int \sqrt{(\sqrt{2})^2 - t^2} dt$$

Rest is easy and work out.

$$④ \int e^{2z} \sqrt{e^{4z} + 6} dz$$

$$= \int e^{2z} \sqrt{(e^{2z})^2 + (\sqrt{6})^2} dz$$

$$\text{put } e^{2z} = v \Rightarrow 2e^{2z} dz = dv$$

$$= \frac{1}{2} \int \sqrt{v^2 + (\sqrt{6})^2} dv$$

Use formula to get answer.

$$\textcircled{5} \int \sec^2 \theta \sqrt{\sec^2 \theta + 3} d\theta$$

$$= \int \sec^2 \theta \sqrt{1 + \tan^2 \theta + 3} d\theta$$

$$= \int \sec^2 \theta \sqrt{\tan^2 \theta + 4} d\theta$$

put $\tan \theta = v$

$$\Rightarrow \sec^2 \theta d\theta = dv$$

$$= \int \sqrt{v^2 + 2^2} dv$$

rest is easy.

$$\textcircled{6} \int \sqrt{x^2 - 4x + 2} dx$$

$$= \int \sqrt{x^2 - 4x + 4 - 2} dx$$

$$= \int \sqrt{(x-2)^2 - (\sqrt{2})^2} dx$$

put $x-2 = t \Rightarrow dx = dt$

$$= \int \sqrt{t^2 - (\sqrt{2})^2} dt$$

rest is easy.

(7) $\int e^{3x} \cos 2x \, dx$ (8) $\int e^{2x} \sin x \, dx$.
 Both you should work out. We have already
 solved general questions $\int e^{ax} \cos(bx) \, dx$ and $\int e^{ax} \sin(bx) \, dx$.

(9) $\int (2x^2+1) e^{x^2} \, dx$

put $x^2 = t$

$\Rightarrow 2x \, dx = dt$

$\Rightarrow dx = \frac{dt}{2\sqrt{t}}$

$= \int (t+1) e^t \frac{dt}{2\sqrt{t}}$

$= \int e^t \left(\sqrt{t} + \frac{1}{2\sqrt{t}} \right) dt$

we know $\frac{d}{dt}(\sqrt{t}) = \frac{1}{2\sqrt{t}}$

$\therefore \int e^t \left(\sqrt{t} + \frac{1}{2\sqrt{t}} \right) dt = e^t \sqrt{t} + K$
 $= e^{x^2} (x) + K$

put $\ln x = t \Rightarrow x = e^t$
 $\Rightarrow dx = e^t \, dt$

(10) $\int \left[\frac{1}{\ln x} - \frac{1}{(\ln x)^2} \right] dx$

$\int \left(\frac{1}{t} - \frac{1}{t^2} \right) e^t \, dt = \frac{e^t}{t} + C = \frac{x}{\ln x} + C$

$$Q \rightarrow \int \sin(\ln x) dx$$

$$\text{put } \ln x = t \Rightarrow x = e^t \Rightarrow dx = e^t dt$$

$$= \int \sin(t) \cdot e^t dt$$

Now use standard by parts technique.

$$A: \rightarrow \int \sin x \ln(\operatorname{cosec} x - \cot x) dx$$

$$= \ln(\operatorname{cosec} x - \cot x) \int \sin x dx - \int \left[\frac{d}{dx} (\ln(\operatorname{cosec} x - \cot x)) \int \sin x dx \right] dx$$

$$= \ln(\operatorname{cosec} x - \cot x) (-\cos x) - \int \left(\frac{\operatorname{cosec}^2 x - \operatorname{cosec} x \cot x}{\operatorname{cosec} x - \cot x} \right) (-\cos x) dx$$

$$= -\cos x \ln(\operatorname{cosec} x - \cot x) + \int (\cos x) \left(\frac{\operatorname{cosec} x (\operatorname{cosec} x - \cot x)}{\operatorname{cosec} x - \cot x} \right) dx$$

$$= -\cos x \ln(\operatorname{cosec} x - \cot x) + \int \frac{\cos x}{\sin x} dx$$

$$= -\cos x \ln(\operatorname{cosec} x - \cot x) + \ln \sin x + \leftarrow$$

Definite Integral

①

Fundamental theorem of Integral Calculus

If $f(x)$ is continuous on $[a, b]$ and $F(x) = \int f(x) dx$

then $\int_a^b f(x) dx = F(b) - F(a)$.

Proof: $\rightarrow \int_a^b f(x) dx = [F(x) + K]_a^b$
 $= (F(b) + K) - (F(a) + K)$
 $= F(b) - F(a)$.

$$\int_a^b \{g(x) \pm h(x)\} dx = \int_a^b g(x) dx \pm \int_a^b h(x) dx$$

$$\int_a^b \lambda g(x) dx = \lambda \int_a^b g(x) dx.$$

e.g ① $\int_1^2 x^3 dx = \left(\frac{x^4}{4}\right)_1^2 = \frac{1}{4}(2^4 - 1^4) = \frac{15}{4}$.

② $\int_0^{\pi/2} \sin x dx = -(\cos x)_0^{\pi/2} = -(\cos \pi/2 - \cos 0)$
 $= -(0 - 1) = 1$.

③ $\int_0^1 \frac{dx}{1+x^2} = \tan^{-1}(x) \Big|_0^1 = \tan^{-1}(1) - \tan^{-1}(0)$
 $= \frac{\pi}{4} - 0 = \frac{\pi}{4}$.

④ $\int_0^{\pi/2} x \cos x dx = \left[x \int \cos x dx - \int \left\{ \frac{d}{dx}(x) \int \cos x dx \right\} dx \right]_0^{\pi/2}$
 $= \left[x(\sin x) - \int (1) \cdot \sin x dx \right]_0^{\pi/2}$
 $= \left\{ x(\sin x) - (-\cos x) \right\}_0^{\pi/2}$

$$= \left[\left\{ \frac{\pi}{2} \sin\left(\frac{\pi}{2}\right) + \cos\left(\frac{\pi}{2}\right) \right\} - \left\{ 0 + \cos(0) \right\} \right]$$

$$= \left(\frac{\pi}{2} + 0 \right) - (0 + 1)$$

$$= \frac{\pi}{2} - 1$$

⑤ $\int_0^{\pi/2} (3x^2 + 2x + \cos x) dx \rightarrow$ Easy left as exercise.

⑥ $\int_2^3 2x e^{x^2} dx$

put $x^2 = t$

$\Rightarrow 2x dx = dt$

$\int_4^9 e^t dt$

$\left(e^t \right)_4^9 = e^9 - e^4$

⑦ $\int_0^{\pi/4} \sin^5 x \cos x dx = \int_0^{\frac{1}{\sqrt{2}}} z^5 dz \quad (\sin x = z)$

$$= \left(\frac{z^6}{6} \right)_0^{\frac{1}{\sqrt{2}}}$$

$$= \frac{1}{6} \left[\left(\frac{1}{\sqrt{2}} \right)^6 \right]$$

$$= \frac{1}{6} \left[\frac{1}{8} \right] = \frac{1}{48}$$

Elementary properties of definite Integral

(2)

$$(i) \int_a^b f(x) dx = - \int_b^a f(x) dx.$$

Proof \rightarrow Let $F'(x) = f(x)$

$$\int_a^b f(x) dx = [F(x)]_a^b = F(b) - F(a).$$

$$\int_b^a f(x) dx = [F(x)]_b^a = F(a) - F(b)$$

$$\therefore \int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$(ii) \int_a^b f(x) dx = \int_a^b f(y) dy = \int_a^b f(z) dz$$

Proof \rightarrow easy. \because definite Integral is independent of symbol of variable of integration

$$(iii) \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx \quad a < c < b.$$

Proof \rightarrow

$$\begin{aligned} \int_a^c f(x) dx + \int_c^b f(x) dx &= |F(x)|_a^c + |F(x)|_c^b \\ &= F(c) - F(a) + F(b) - F(c) \\ &= F(b) - F(a) \\ &= |F(x)|_a^b \\ &= \int_a^b f(x) dx. \end{aligned}$$

2

$$\textcircled{1} \int_1^4 [x] dx$$

$$= \int_1^2 [x] dx + \int_2^3 [x] dx + \int_3^4 [x] dx$$

$$= \int_1^2 1 dx + \int_2^3 2 dx + \int_3^4 3 dx$$

$$= (2-1) + 2(3-2) + 3(4-3)$$

$$= 1 + 2 + 3 = 6.$$

$$\textcircled{2} \int_{-3}^4 |x| dx$$

$$= \int_{-3}^0 |x| dx + \int_0^4 |x| dx$$

$$= -\int_{-3}^0 x dx + \int_0^4 x dx$$

$$= -\left(\frac{x^2}{2}\right)_{-3}^0 + \left(\frac{x^2}{2}\right)_0^4$$

$$= -\frac{1}{2}(0-9) + \frac{1}{2}(16-0)$$

$$= \frac{9}{2} + \frac{16}{2} = \frac{25}{2}.$$

$$\textcircled{3} \int_0^1 (2x+1)^4 dx$$

$$= \frac{(2x+1)^5}{\frac{10}{5}} \Big|_0^1$$

$$= \frac{1}{10} [3^5 - 1^5] = \frac{242}{10} = \frac{121}{5}.$$

(5)

$$\int_0^1 x^7 (1+x^8)^{1/3} dx.$$

put $1+x^8 = t$ $t=1$ when $x=0$
 $t=2$ when $x=1$

$$\Rightarrow 8x^7 = \frac{dt}{dx}$$

$$\Rightarrow x^7 dx = \frac{dt}{8}$$

$$\frac{1}{8} \int_1^2 dt t^{1/3}$$

$$= \frac{1}{8} \left[\frac{t^{4/3}}{4/3} \right]_1^2$$

$$= \frac{1}{8} \left[\frac{3}{4} t^{4/3} \right]_1^2$$

$$= \frac{3}{32} (2^{4/3} - 1).$$

(6)

$$\int_0^{\pi/2} (\cos x - \sin x) dx$$

$$= \int_0^{\pi/2} \cos x dx - \int_0^{\pi/2} \sin x dx$$

$$= \left[\sin x \right]_0^{\pi/2} - \left[-\cos x \right]_0^{\pi/2}$$

$$= [\sin(\pi/2) - \sin 0] + [\cos(\pi/2) - \cos 0]$$

$$= (1-0) + (0-1) = 0.$$

$$\begin{aligned}
 (7) \quad & \int_{\pi/6}^{\pi/3} \frac{1 + \cos 2x}{1 - \cos 2x} dx \\
 &= \int_{\pi/6}^{\pi/3} \frac{\cancel{2} \cos^2 x}{\cancel{2} \sin^2 x} dx \\
 &= \int_{\pi/6}^{\pi/3} \cot^2 x dx \\
 &= -\operatorname{cosec}(x) \Big|_{\pi/6}^{\pi/3} \\
 &= -\left[\operatorname{cosec}\left(\frac{\pi}{3}\right) - \operatorname{cosec}\left(\frac{\pi}{6}\right) \right] \\
 &= -\left[\frac{2}{\sqrt{3}} - 2 \right] \\
 &= 2 - \frac{2}{\sqrt{3}}
 \end{aligned}$$

$$\begin{aligned}
 (8) \quad & \int_0^{3/2} [2x] dx \\
 &= \int_0^{1/2} [2x] dx + \int_{1/2}^1 [2x] dx + \int_1^{3/2} [2x] dx \\
 & 0 \leq x < \frac{1}{2} \Rightarrow 0 \leq 2x < 1 \Rightarrow [2x] = 0 \\
 & \frac{1}{2} \leq x < 1 \Rightarrow 1 \leq 2x < 2 \Rightarrow [2x] = 1 \\
 & 1 \leq x < \frac{3}{2} \Rightarrow 2 \leq 2x < 3 \Rightarrow [2x] = 2 \\
 &= \int_0^{1/2} 0 dx + \int_{1/2}^1 1 dx + 2 \int_1^{3/2} dx \\
 &= \left(1 - \frac{1}{2}\right) + 2\left(\frac{3}{2} - 1\right) = \frac{1}{2} + 2\left(\frac{1}{2}\right) = 1 + \frac{1}{2} \\
 &= \frac{3}{2}
 \end{aligned}$$

Some More properties of definite Integral (4)

$$\textcircled{1} \int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

In particular $\int_0^a f(x) dx = \int_0^a f(a-x) dx$.

$$\textcircled{2} \int_{-a}^a f(x) dx = \begin{cases} 2 \int_0^a f(x) dx & \text{if } f(-x) = f(x) \\ 0 & \text{if } f(-x) = -f(x) \end{cases}$$

$$\textcircled{3} \int_0^{2a} f(x) dx = \begin{cases} 2 \int_0^a f(x) dx & \text{if } f(2a-x) = f(x) \\ 0 & \text{if } f(2a-x) = -f(x) \end{cases}$$

$$\text{Q:} \rightarrow \int_1^2 \frac{\sqrt{x} dx}{\sqrt{3-x} + \sqrt{x}}$$

$$\text{Let } I = \int_1^2 \frac{\sqrt{x} dx}{\sqrt{3-x} + \sqrt{x}} = \int_1^2 \frac{\sqrt{3-x}}{\sqrt{x} + \sqrt{3-x}} dx \text{ using } \textcircled{1}$$

$$\therefore 2I = \int_1^2 \left(\frac{\sqrt{x} + \sqrt{3-x}}{\sqrt{x} + \sqrt{3-x}} \right) dx$$

$$= \int_1^2 dx = (x)_1^2 = 2 - 1 = 1$$

$$\therefore \boxed{I = \frac{1}{2}}$$

$$Q: \rightarrow \int_0^a \frac{dx}{x + \sqrt{a^2 - x^2}}$$

put $x = a \sin \theta$ $x=0 \Rightarrow \theta=0, x=a \Rightarrow \theta = \pi/2$

$$\Rightarrow dx = a \cos \theta d\theta$$

$$= \int_0^{\pi/2} \frac{a \cos \theta d\theta}{a \sin \theta + \sqrt{a^2 - a^2 \sin^2 \theta}}$$

$$= \int_0^{\pi/2} \frac{a \cos \theta d\theta}{a \sin \theta + \sqrt{a^2 \cos^2 \theta}}$$

$$= \int_0^{\pi/2} \frac{a \cos \theta d\theta}{a \sin \theta + a \cos \theta}$$

$$= \int_0^{\pi/2} \frac{\sin \theta \cos \theta d\theta}{\sin \theta + \cos \theta}$$

$$= \int_0^{\pi/2} \frac{\cos(\pi/2 - \theta) d\theta}{\sin(\pi/2 - \theta) + \cos(\pi/2 - \theta)}$$

$$= \int_0^{\pi/2} \frac{\sin \theta d\theta}{\cos \theta + \sin \theta}$$

$$2I = \int_0^{\pi/2} \left(\frac{\sin \theta + \cos \theta}{\sin \theta + \cos \theta} \right) d\theta$$

$$= \int_0^{\pi/2} d\theta = (\theta)_0^{\pi/2} = \pi/2$$

$$\boxed{I = \pi/4}$$

(5)

$$Q: \rightarrow \int_0^{\pi/2} \frac{\sin \theta d\theta}{\sin \theta + \cos \theta} = \int_0^{\pi/2} \frac{\cos \theta d\theta}{\sin \theta + \cos \theta} \quad (\text{Imp})$$

$$= \int_0^{\pi/2} \frac{d\theta}{1 + \cot \theta} = \int_0^{\pi/2} \frac{d\theta}{1 + \tan \theta} = \pi/4$$

All exercises are same and proofs available in the last exercise solved.

$$Q: \rightarrow \int_0^{\pi/2} \frac{\sqrt{\sin \theta} d\theta}{\sqrt{\sin \theta} + \sqrt{\cos \theta}} = \int_0^{\pi/2} \frac{\sqrt{\cos \theta} d\theta}{\sqrt{\cos \theta} + \sqrt{\sin \theta}} = \int_0^{\pi/2} \frac{d\theta}{1 + \sqrt{\tan \theta}} = \int_0^a \frac{dx}{1 + \sqrt{x}}$$

Use property $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

Ans: $\rightarrow \pi/4$

$$Q: \rightarrow \int_0^{\pi/4} \log(1 + \tan \theta) d\theta \quad (\text{Imp})$$

$$I = \int_0^{\pi/4} \ln(1 + \tan \theta) d\theta = \int_0^{\pi/4} \ln[1 + \tan(\pi/4 - \theta)] d\theta$$

$$= \int_0^{\pi/4} \ln \left[1 + \frac{1 - \tan \theta}{1 + \tan \theta} \right] d\theta = \int_0^{\pi/4} \ln \left(\frac{1 + \tan \theta + 1 - \tan \theta}{1 + \tan \theta} \right) d\theta$$

$$= \int_0^{\pi/4} \ln \left(\frac{2}{1 + \tan \theta} \right) d\theta$$

$$= \int_0^{\pi/4} \{ \ln(2) - \ln(1 + \tan \theta) \} d\theta$$

$$= \ln(2) \int_0^{\pi/4} d\theta - \int_0^{\pi/4} \ln(1 + \tan \theta) d\theta$$

$$= \pi/4 \ln 2 - I \quad \Rightarrow 2I = \pi/4 \ln 2$$

$$\Rightarrow I = \frac{\pi}{8} \ln 2$$

Q: $\rightarrow \int_0^{\pi/2} \frac{\sin \theta d\theta}{\sin \theta + \cos \theta} = \int_0^{\pi/2} \frac{\cos \theta d\theta}{\sin \theta + \cos \theta}$ (Imp)

$= \int_0^{\pi/2} \frac{d\theta}{1 + \cot \theta} = \int_0^{\pi/2} \frac{d\theta}{1 + \tan \theta} = \pi/4$

All exercises are same and prefix variable in the last exercise solved.

Q: $\rightarrow \int_0^{\pi/2} \frac{\sqrt{\sin \theta} d\theta}{\sqrt{\sin \theta} + \sqrt{\cos \theta}} = \int_0^{\pi/2} \frac{\sqrt{\cos \theta} d\theta}{\sqrt{\cos \theta} + \sqrt{\sin \theta}} = \int_0^{\pi/2} \frac{d\theta}{1 + \sqrt{\tan \theta}} = \int_0^a \frac{d\theta}{1 + \sqrt{\tan \theta}}$

Ans: $\rightarrow \pi/4$ UAC property $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

Q: $\rightarrow \int_0^{\pi/4} \log(1 + \tan \theta) d\theta$ (Imp)

$I = \int_0^{\pi/4} \ln(1 + \tan \theta) d\theta = \int_0^{\pi/4} \ln[1 + \tan(\pi/4 - \theta)] d\theta$

$= \int_0^{\pi/4} \ln \left[1 + \frac{1 - \tan \theta}{1 + \tan \theta} \right] d\theta = \int_0^{\pi/4} \ln \left(\frac{1 + \tan \theta + 1 - \tan \theta}{1 + \tan \theta} \right) d\theta$

$= \int_0^{\pi/4} \ln \left(\frac{2}{1 + \tan \theta} \right) d\theta$

$= \int_0^{\pi/4} \{ \ln(2) - \ln(1 + \tan \theta) \} d\theta$

$= \ln(2) \int_0^{\pi/4} d\theta - \int_0^{\pi/4} \ln(1 + \tan \theta) d\theta$

$= \pi/4 \ln 2 - I \Rightarrow 2I = \pi/4 \ln 2 \Rightarrow I = \frac{\pi}{8} \ln 2$

Q: $\rightarrow \int_0^{\pi/2} \frac{\cos x - \sin x}{1 + \sin x \cos x} dx = 0$ (prove).

Solⁿ: $\rightarrow f(x) = \frac{\cos x - \sin x}{1 + \sin x \cos x}$
 $f(\frac{\pi}{2} - x) = \frac{\cos(\frac{\pi}{2} - x) - \sin(\frac{\pi}{2} - x)}{1 + \sin(\frac{\pi}{2} - x) \cos(\frac{\pi}{2} - x)}$
 $= \frac{\sin x - \cos x}{1 + \sin x \cos x} = -f(x)$

$\therefore \int_0^{\pi/2} f(x) dx = 0$ ($\because \int_0^{2a} f(x) dx = 0$ if $f(2a-x) = -f(x)$)

Q: $\rightarrow \int_0^{\pi} \sin x \cos^2 x dx = 2 \int_0^{\pi/2} \cos^2 x \sin x dx$ ($\because f(\pi-x) = f(x)$)
 $= -2 \int_1^0 t^2 dt$ put $\cos x = t$
 $\Rightarrow -\sin x dx = dt$
 $= 2 \int_0^1 t^2 dt = 2 \left(\frac{t^3}{3} \right)_0^1 = \frac{2}{3} (1-0) = \frac{2}{3}$

Q: $\rightarrow \int_1^3 \frac{\sqrt{x} dx}{\sqrt{4-x} + \sqrt{x}}$

Solⁿ: $\rightarrow I = \int_1^3 \frac{\sqrt{x} dx}{\sqrt{4-x} + \sqrt{x}} = \int_1^3 \frac{\sqrt{1+3-x} dx}{\sqrt{4-(1+3-x)} + \sqrt{1+3-x}}$

$= \int_1^3 \frac{\sqrt{4-x} dx}{\sqrt{x} + \sqrt{4-x}} = I$

$$\therefore 2I = \int_1^3 \left(\frac{\sqrt{x} + \sqrt{4-x}}{\sqrt{4-x} + \sqrt{x}} \right) dx$$

$$= \int_1^3 dx = (x)_1^3 = 3-1 = 2.$$

$$\Rightarrow \boxed{I=1}$$

$$Q: \rightarrow \int_{\pi/6}^{\pi/3} \frac{dx}{1 + \sqrt{\cot x}}$$

$$I = \int_{\pi/6}^{\pi/3} \frac{dx}{1 + \frac{\sqrt{\cos x}}{\sqrt{\sin x}}}$$

$$= \int_{\pi/6}^{\pi/3} \frac{\sqrt{\sin x} dx}{\sqrt{\sin x} + \sqrt{\cos x}}$$

$$= \int_{\pi/6}^{\pi/3} \frac{dx}{\sqrt{\sin(\frac{\pi}{3} + \frac{\pi}{6} - x)} + \sqrt{\cos(\frac{\pi}{3} + \frac{\pi}{6} - x)}}$$

$$= \int_{\pi/6}^{\pi/3} \frac{dx}{\sqrt{\sin(\frac{\pi}{2} - x)} + \sqrt{\cos(\frac{\pi}{2} - x)}}$$

$$= \int_{\pi/6}^{\pi/3} \frac{\sqrt{\cos x} dx}{\sqrt{\sin x} + \sqrt{\cos x}} = I$$

$$\therefore 2I = \int_{\pi/6}^{\pi/3} \left(\frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} \right) dx = \int_{\pi/6}^{\pi/3} dx$$

$$= \frac{\pi}{3} - \frac{\pi}{6} = \frac{\pi}{6}$$

$$\therefore \boxed{I = \frac{\pi}{12}}$$

$$\begin{aligned}
 Q: \rightarrow & \int_0^1 x(1-x)^{100} dx \\
 &= \int_0^1 (0+1-x)(1-(1-x))^{100} dx \\
 &= \int_0^1 (1-x)(x^{100}) dx \\
 &= \int_0^1 (x^{100} - x^{101}) dx \\
 &= \int_0^1 x^{100} dx - \int_0^1 x^{101} dx \\
 &= \left[\frac{x^{101}}{101} \right]_0^1 - \left[\frac{x^{102}}{102} \right]_0^1 \\
 &= \left(\frac{1}{101} - 0 \right) - \left(\frac{1}{102} - 0 \right) \\
 &= \frac{1}{101} - \frac{1}{102} = \frac{102-101}{(101)(102)} = \frac{1}{(101)(102)}
 \end{aligned}$$

$$Q: \rightarrow \int_0^{\pi/2} \ln(\sin x) dx = \int_0^{\pi/2} \ln(\cos x) dx = \frac{\pi}{2} \ln\left(\frac{1}{2}\right)$$

$$\begin{aligned}
 \text{Sol}^n: \rightarrow I &= \int_0^{\pi/2} \ln(\sin x) dx = \int_0^{\pi/2} \ln(\sin(\frac{\pi}{2}-x)) dx \\
 &= \int_0^{\pi/2} \ln(\cos x) dx = I
 \end{aligned}$$

$$2I = \int_0^{\pi/2} \{ \ln(\sin x) + \ln(\cos x) \} dx$$

$$\begin{aligned}
&= \int_0^{\pi/2} \ln(\sin x \cos x) dx \\
&= \int_0^{\pi/2} \ln\left(\frac{2 \sin x \cos x}{2}\right) dx \\
&= \int_0^{\pi/2} \ln\left(\frac{\sin 2x}{2}\right) dx \\
&= \int_0^{\pi/2} \ln(\sin 2x) dx - \int_0^{\pi/2} (\ln 2) \cdot dx \\
&= \int_0^{\pi/2} \ln(\sin 2x) dx - (\ln 2) \cdot \frac{\pi}{2}
\end{aligned}$$

put $2x = v$

$$\Rightarrow 2 dx = dv$$

$$\Rightarrow dx = \frac{dv}{2}$$

$$= \frac{1}{2} \int_0^{\pi} \ln(\sin v) dv - \frac{\pi}{2} (\ln 2)$$

$$= \left(\frac{1}{2}\right) \cdot \cancel{2} \int_0^{\pi/2} \ln(\sin x) dx - \frac{\pi}{2} \ln(2)$$


$$\Rightarrow 2I = I - \frac{\pi}{2} \ln 2$$

$$\Rightarrow I = -\frac{\pi}{2} \ln 2 \Rightarrow I = \frac{\pi}{2} \ln 2^{-1} = \frac{\pi}{2} \ln\left(\frac{1}{2}\right)$$


Area Under plane curves

(1)

$\int_a^b f(x) dx$ represents the area under $y = f(x)$ above x -axis between ordinates $x = a$ and $x = b$.



Area betⁿ $x = f(y)$, $x = 0$ and the abscissae $y = c$, $y = d$ is equal to $\int_c^d f(y) dy$




Area betⁿ 2 curves $y = f(x)$, $y = g(x)$

with $g(x) < f(x)$ in $[a, b]$ is given by


$$A = \int_a^b \{f(x) - g(x)\} dx.$$

Graph of standard curves


circle: $\rightarrow x^2 + y^2 = r^2$



circle $\rightarrow (x-h)^2 + (y-k)^2 = r^2$




circle $(x-h)^2 + (y-k)^2 = a^2$




Ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$




Parabola

$$y^2 = 4ax \quad (a > 0)$$



$$x^2 = 4ay \quad (a > 0)$$




Q: → Find the area of circle $x^2 + y^2 = a^2$

$$A = 4 \int_0^a (\sqrt{a^2 - x^2}) dx$$

$$= 4 \left[\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) \right]_0^a$$

$$= 4 \left[\frac{a^2}{2} \cdot \sin^{-1}(1) \right] = 4 \left[\frac{a^2}{2} \cdot \frac{\pi}{2} \right] = \pi a^2$$



Q: → Find the area of ellipse. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$$A = 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx$$


$$= 4 \frac{b}{a} \int_0^a \sqrt{a^2 - x^2} dx$$

$$= \frac{4b}{a} \left[\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) \right]_0^a$$

$$= \frac{4b}{a} \left[\left\{ \frac{a}{2} \sqrt{a^2 - a^2} + \frac{a^2}{2} \sin^{-1}(1) \right\} - \left\{ 0 + 0 \right\} \right]$$

$$= \frac{4b}{a} \cdot \frac{a^2}{2} \cdot \frac{\pi}{2}$$

$$= \pi ab$$




Q: → Find area bounded by

$$y = e^x, y = 0, x = 2, x = 4$$

$$\text{Ans: } \rightarrow A = \int_2^4 y dx = \int_2^4 e^x dx = (e^x)_2^4 = e^4 - e^2$$

$$y = x^2, y = 0, x = 1$$

$$A = \int_0^1 x^2 dx = \left. \frac{x^3}{3} \right|_0^1 = \frac{1}{3}$$



(3)

Q: $\rightarrow y = \sin x, y=0, x = \pi/2.$

A: $\rightarrow A = \int_0^{\pi/2} \sin x \, dx = (-\cos x)_0^{\pi/2}$
 $= -(0 - 1) = 1.$

Q: $\rightarrow xy = a^2, y=0, x = \alpha, x = \beta \ (\beta > \alpha > 0).$

A: $\rightarrow \text{Area} = \int_{\alpha}^{\beta} y \, dx = \int_{\alpha}^{\beta} \left(\frac{a^2}{x}\right) \, dx = a^2 \ln x \Big|_{\alpha}^{\beta}$
 $= a^2 (\ln b - \ln a)$
 $= a^2 \ln\left(\frac{b}{a}\right).$



Find area enclosed by $x=0, y=2, y=3.$

① $y = e^x, x=0, y=2, y=3.$

② $y^2 = x, x=0, y=1$

③ $xy = a^2, x=0, y = \alpha, y = \beta \ (0 < \alpha < \beta)$

④ $y^2 = x^3, x=0, y=1$

Solⁿ of ②


Area = $\int_0^1 x \, dy = \int_0^1 y^2 \, dy$
 $= \frac{y^3}{3} \Big|_0^1$
 $= \frac{1}{3}$

Rest you should work out.

Q: \rightarrow Find area of parabola $y^2 = 4ax$ bounded

by $x = a$.

Sol: \rightarrow



$$\text{Area} = 2 \int_0^a y \, dx$$

$$= 2 \int_0^a \sqrt{4ax} \, dx$$

$$= 2 \int_0^a (2)\sqrt{a} \sqrt{x} \, dx$$


$$= (4)(\sqrt{a}) \left[\frac{x^{3/2}}{3/2} \right]_0^a$$

$$= \frac{8}{3} (a)^{1/2} \cdot (a)^{3/2}$$

$$= \frac{8}{3} (a)^2$$

Q: \rightarrow Find area common to two parabolas

$y^2 = 4ax$ and $x^2 = 4ay$



Sol: \rightarrow Area = \int

$$y^2 = 4ax \Rightarrow x = \frac{y^2}{4a}$$

$$\therefore \frac{y^4}{16a^2} = 4ay \Rightarrow y^3 = 64a^3$$

$$\Rightarrow y = 4a$$

$$\therefore x = \frac{16a^2}{4a} = 4a$$

1

$$\begin{aligned}
\therefore \text{Area} &= \int_0^{4a} (y_1 - y_2) dx \\
&= \int_0^{4a} \left(\sqrt{4ax} - \frac{x^2}{4a} \right) dx \\
&= \int_0^{4a} \left(2a^{1/2} x^{1/2} - \frac{1}{4a} x^2 \right) dx \\
&= 2\sqrt{a} \left(\frac{x^{3/2}}{3/2} \right) - \frac{1}{4a} \cdot \frac{x^3}{3} \Bigg|_0^{4a} \\
&= \left[\frac{4a^{1/2}}{3} x^{3/2} - \frac{1}{12a} x^3 \right]_0^{4a} \\
&= \frac{4a^{1/2}}{3} \cdot (4a)^{3/2} - \frac{1}{12a} (4a)^3 \\
&= \frac{4}{3} a^{1/2} (8) a^{3/2} - \frac{1}{12a} \cdot 64a^3 \\
&= \frac{32}{3} a^2 - \frac{16}{3} a^2 = \frac{16a^2}{3}
\end{aligned}$$

Ex Find the area betⁿ parabolas $y^2 = x$ and $x^2 = y$